

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
9 February 2006 (09.02.2006)

PCT

(10) International Publication Number
WO 2006/014359 A2

(51) International Patent Classification:
C07D 401/00 (2006.01)

(21) International Application Number:
PCT/US2005/023554

(22) International Filing Date: 1 July 2005 (01.07.2005)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/585,292 2 July 2004 (02.07.2004) US

(71) Applicant (for all designated States except US): ICOS CORPORATION [US/US]; 22021 20th Avenue S.E., Bothell, WA 98021 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): GAUDINO, John, Joseph [US/US]; 4224 Prairie Fire Circle, Longmont, CO 80503 (US). COOK, Adam, Wade [US/US]; 5983 Star View Drive, Broomfield, CO 80020 (US).

(74) Agent: NAPOLI, James, J.; Marshall, Gerstein & Borun LLP, 233 S. Wacker Drive, Suite 6300, Sears Tower, Chicago, IL 60606-6357 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: COMPOUNDS USEFUL FOR INHIBITING CHK1

(57) Abstract: Substituted urea compounds useful in the treatment of diseases and conditions related to DNA damage or lesions in DNA replication are disclosed. Methods of making the compounds, and their use as therapeutic agents, for example, in treating cancer and other diseases characterized by defects in DNA replication, chromosome segregation, or cell division also are disclosed.

BEST AVAILABLE COPY

WO 2006/014359 A2

COMPOUNDS USEFUL FOR INHIBITING CHK1**CROSS-REFERENCE TO RELATED APPLICATION**

This application claims the benefit of U.S.
provisional application Serial No. 60/585,292, filed July
5 2, 2004.

FIELD OF THE INVENTION

The present invention relates to compounds
useful for inhibiting enzymes that maintain and repair
the integrity of genetic material. More particularly,
10 the present invention relates to a series of aryl- and
heteroaryl-substituted urea compounds, methods of making
the compounds, and their use as therapeutic agents, for
example, in treating cancer and other diseases character-
ized by defects in deoxyribonucleic acid (DNA) replica-
15 tion, chromosome segregation, or cell division.

BACKGROUND OF THE INVENTION

A large variety of diseases, conditions, and
disorders (hereinafter "indications") are characterized
as involving aberrantly proliferating cells. As used
20 herein, "aberrantly proliferating cells" (or "aberrant
cell proliferation") means cell proliferation that
deviates from the normal, proper, or expected course.
For example, aberrant cell proliferation includes in-
appropriate proliferation of cells wherein DNA or other
25 cellular components have become damaged or defective.
Aberrant cell proliferation also includes indications
caused by, mediated by, or resulting in inappropriately
high levels of cell division, inappropriately low levels
of cell death (e.g., apoptosis), or both. Such indica-

tions can be characterized, for example, by single or multiple local abnormal proliferations of cells, groups of cells or tissue(s), and include cancerous (benign or malignant) and noncancerous indications.

5 By definition, all cancers (benign and malignant) involve some form of aberrant cell proliferation. Some noncancerous indications also involve aberrant cell proliferation. Examples of noncancerous indications involving aberrant cell proliferation include rheumatoid
10 arthritis, psoriasis, vitiligo, Wegener's granulomatosis, and systemic lupus.

 One approach to treating indications involving aberrantly proliferating cells involves the use of DNA damaging agents. These agents are designed to kill
15 aberrantly proliferating cells by disrupting vital cellular processes such as DNA metabolism, DNA synthesis, DNA transcription, and microtubule spindle formation. They also can operate, for example, by introducing lesions into DNA that perturb chromosomal structural integrity.
20 DNA damaging agents are designed and administered in ways that attempt to induce maximum damage and consequent cell death in aberrantly proliferating cells with a minimum damage to normal, healthy cells.

 A large variety of DNA damaging agents have
25 been developed to date, including chemotherapeutics and radiation, and others are in development. Unfortunately, the effectiveness of DNA damaging agents in treating conditions involving aberrant cell proliferation have been less than desired, particularly in the treatment of
30 cancer. The selectivity of such agents for aberrantly proliferating cells over healthy cells (sometimes referred to as the therapeutic index) often is marginal.

 Moreover, all cells have sensing and repair mechanisms that can work at cross purposes to DNA

damaging agents. Such sensing mechanisms, called cell cycle checkpoints, help to maintain the order of the various cell replication stages and to ensure that each step is executed with high fidelity (Hartwell et al.,
5 *Science*, 246:629-34 (1989); Weinert et al., *Genes Dev.*, 8:652 (1994)). When cells detect DNA damage, including damage purposefully induced by DNA damaging agents, certain signaling pathways activate cell cycle checkpoints and the cell replication cycle temporarily ceases
10 ("arrests"). This arrest allows cells time to repair their DNA, often to a degree sufficient to allow the affected cells to continue to survive and proliferate. In the case of aberrantly proliferating cells, this repair is unwanted, as it may undermine efforts to induce
15 DNA damage sufficient to kill such cells.

For example, the chemotherapeutic agent called GEMZAR™ (gemcitabine, or 2',2' difluoro-2'-deoxycytidine) damages DNA by incorporating itself into DNA during synthesis. Left unrepaired, damaged DNA generally is rendered incapable of sustaining life. In many targeted
20 cells, however, cell cycle checkpoints detect the improperly made (or otherwise damaged) DNA. The activated cell cycle checkpoints trigger cell cycle arrest for a time sufficient to allow damaged DNA to be repaired. This is
25 one way in which aberrantly proliferating cells are theorized to resist the cell-killing effect of DNA-damaging agents such as chemotherapeutics, radiation, and other therapies.

Other DNA-damaging agents cause tumor cells to
30 arrest in S-phase. Tumor cells have been observed to resist certain chemotherapeutics simply by arresting in S phase while the chemotherapeutic agent is being administered. Then, as soon as the drug is removed, DNA damage is repaired, cell cycle arrest ceases, and the cells

progress through the remainder of the cell cycle (Shi et al., *Cancer Res.* 61:1065-72, 2001). Other therapeutics cause cell cycle arrest at other checkpoints, including G1 and G2. Inhibition of various DNA damage checkpoints
5 therefore is expected to assist in preventing cells from repairing therapeutically induced DNA damage and to sensitize targeted cells to DNA damaging agents. Such sensitization is in turn expected to increase the therapeutic index of these therapies.

10 The cell cycle is structurally and functionally the same in its basic process and mode of regulation across all eukaryotic species. The mitotic (somatic) cell cycle consists of four phases: the G1 (gap) phase, the S (synthesis) phase, the G2 (gap) phase, and the M
15 (mitosis) phase. The G1, S, and G2 phases are collectively referred to as interphase of the cell cycle. During the G1 phase, biosynthetic activities of the cell progress at a high rate. The S phase begins when DNA synthesis starts, and ends when the DNA content of the
20 nucleus of the cell has been replicated and two identical sets of chromosomes are formed.

 The cell then enters the G2 phase, which continues until mitosis starts. In mitosis, the chromosomes pair and separate, two new nuclei form, and cytokinesis
25 occurs in which the cell splits into two daughter cells each receiving one nucleus containing one of the two sets of chromosomes. Cytokinesis terminates the M phase and marks the beginning of interphase of the next cell cycle. The sequence in which cell cycle events proceed is tightly
30 ly regulated, such that the initiation of one cell cycle event is dependent on the completion of the prior cell cycle event. This allows fidelity in the duplication and segregation of genetic material from one generation of somatic cells to the next.

It has been reported that cell cycle checkpoints comprise at least three distinct classes of polypeptides, which act sequentially in response to cell cycle signals or defects in chromosomal mechanisms (Carr, *Science*, 271:314-15, 1996). The first class is a family of proteins that detect or sense DNA damage or abnormalities in the cell cycle. These sensors include Ataxia-telangiectasia Mutated protein (Atm) and Ataxia-Telangiectasia Rad-related protein (Atr). The second class of polypeptides amplify and transmit the signal detected by the detector and is exemplified by Rad53 (Alen et al., *Genes Dev.* 8:2416-88, 1994) and Chk1. A third class of polypeptides includes cell cycle effectors, such as p53, which mediate a cellular response, for example, arrest of mitosis and apoptosis.

Much of the current understanding of the function of cell cycle checkpoints has been derived from the study of tumor derived cell lines. In many cases, tumor cells have lost key cell cycle check points (Hartwell et al., *Science* 266:1821-28, 1994). It has been reported that a key step in the evolution of cells to a neoplastic state is the acquisition of mutations that inactivate cell cycle checkpoint pathways, such as those involving p53 (Weinberg, *Cell* 81:323-30, 1995; Levine, *Cell* 88:3234-31, 1997). Loss of these cell cycle checkpoints results in the replication of tumor cells despite DNA damage.

Noncancerous tissue, which has intact cell cycle checkpoints, typically is insulated from temporary disruption of a single checkpoint pathway. Tumor cells, however, have defects in pathways controlling cell cycle progression such that the perturbation of additional checkpoints renders them particularly sensitive to DNA damaging agents. For example, tumor cells that contain

mutant p53 are defective both in the G1 DNA damage checkpoint and in the ability to maintain the G2 DNA damage checkpoint (Bunz et al., *Science*, 282:1497-501, 1998). Checkpoint inhibitors that target initiation of the G2 checkpoint or the S phase checkpoint are expected to further cripple the ability of these tumor cells to repair DNA damage and, therefore, are candidates to enhance the therapeutic index of both radiation and systemic chemotherapy (Gesner, Abstract at SRI Conference: Protein Phosphorylation and Drug Discovery World Summit, March 2003).

In the presence of DNA damage or any impediment to DNA replication, the checkpoint proteins Atm and Atr initiate a signal transduction pathway leading to cell cycle arrest. Atm has been shown to play a role in a DNA damage checkpoint in response to ionizing radiation. Atr is stimulated by agents that cause double strand DNA breaks, single strand DNA breaks, and agents that block DNA radiation.

Chk1 is a protein kinase that lies downstream from Atm and/or Atr in the DNA damage checkpoint signal transduction pathway (Sanchez et al., *Science*, 277:1497-501, 1997; U.S. Patent No. 6,218,109). In mammalian cells, Chk1 is phosphorylated in response to agents that cause DNA damage including ionizing radiation, ultraviolet (UV) light, and hydroxyurea (Sanchez et al., supra; Lui et al., *Genes Dev.*, 14:1448-59, 2000). This phosphorylation, which activates Chk1 in mammalian cells, is dependent on Atm (Chen et al., *Oncogene*, 18:249-56, 1999) and Atr (Lui et al., supra). Furthermore, Chk1 has been shown to phosphorylate both weel (O'Connell et al., *EMBO J.*, 16:545-54, 1997) and Pds1 (Sanchez et al., *Science*, 286:1166-71, 1999), gene products known to be important in cell cycle control.

These studies demonstrate that mammalian Chk1 plays a role in the Atm dependent DNA damage checkpoint leading to arrest at S phase. A role for Chk1 in the S phase mammalian cells has recently been elucidated

5 (Feijoo et al., *J. Cell. Biol.*, 154:913-23, 2001; Zhao et al., *Proc. Nat. Acad. Sci. USA*, 99:14795-800, 2002; Xiao et al., *J Biol Chem.*, 278(24):21767-773, 2003; Sorensen et al., *Cancer Cell*, 3(3):247-58, 2003) highlighting the role of Chk1 in monitoring the integrity of DNA synthe-
10 sis. Chk1 invokes an S-phase arrest by phosphorylating Cdc25A, which regulates cyclinA/cdk2 activity (Xiao et al., supra and Sorensen et al., supra). Chk1 also invokes a G2 arrest by phosphorylating and inactivating Cdc25C, the dual specificity phosphatase that normally
15 dephosphorylates cyclin-B/cdc2 (also known as Cdk1) as cells progress from G2 into mitosis (Fernery et al., *Science*, 277:1495-7, 1997; Sanchez et al., supra; Matsuoka et al., *Science*, 282:1893-97, 1998; and Blasina et al., *Curr. Biol.*, 9:1-10, 1999). In both cases,
20 regulation of Cdk activity induces a cell cycle arrest to prevent cells from entering mitosis in the presence of DNA damage or unreplicated DNA.

Additional classes of cell cycle checkpoint inhibitors operate at either the G1 or G2/M phase. UCN-
25 01, or 7-hydroxystaurosporine, originally was isolated as a nonspecific kinase inhibitor having its primary effect on protein kinase C, but recently it has been found to inhibit the activity of Chk1 and abrogate the G2 cell cycle checkpoint (Shi et al., supra). Thus, because UCN-
30 01 is a nonselective Chk1 inhibitor, it is toxic to cells at high doses. At low doses, it nonspecifically inhibits many cellular kinases and also inhibits the G1 checkpoint (Tenzer et al., *Curr. Med. Chem. Anti-Cancer Agents*, 3:35-46, 2003).

UCN-01 has been used in conjunction with cancer therapies, such as radiation, the anti-cancer agent camptothecin (Tenzer et al., supra), and gemcitabine (Shi et al., supra), with limited success. In addition, UCN-01 has been used to potentiate the effects of temozolomide (TMZ)-induced DNA mismatch repair (MMR) in glioblastoma cells (Hirose et al., *Cancer Res.*, 61:5843-49, 2001). In the clinic, UCN-01 is not an effective chemotherapeutic as expected, possibly due to a failure in treatment scheduling and a lack of identification of particular key molecular targets (Grant et al., *Drug Resistance Updates*, 6:15-26, 2003). Thus, Mack et al. report cell cycle-dependent potentiation of cisplatin by UCN-01 in a cultured nonsmall-cell lung carcinoma cell line, but do not identify with specificity the key cell cycle checkpoint(s) targeted by UCN-01. (Mack et al., *Cancer Chemother. Pharmacol.*, 51(4):337-48, 2003).

Several other strategies exist for sensitizing tumor cells to treatment with cell cycle affecting chemotherapeutics. For example, administration of 2-aminopurine abrogates multiple cell cycle checkpoint mechanisms, such as mimosine-induced G1 arrest or hydroxyurea-induced S phase arrest, allowing the cell to progress into and through mitosis (Andreassen et al., *Proc Natl Acad Sci USA*, 86:2272-76, 1992). Caffeine, a methylxanthine, has also been used to enhance cytotoxicity of DNA-damaging agents, such as cisplatin and ionizing radiation, by mediating progression through the G2 checkpoint and thereby inducing cell death. (Bracey et al., *Clin. Cancer Res.*, 3:1371-81, 1997). However, the dose of caffeine used to accomplish the cell cycle abrogation exceeds clinically acceptable levels and is not a viable therapeutic option. Additionally, antisense nucleotides to Chk1 kinase have been used to increase

sensitivity to the topoisomerase inhibitor BNP1350 (Yin et al., *Biochem. Biophys. Res. Commun.*, 295:435-44, 2002), but demonstrate problems typically associated with antisense treatment and gene therapy.

5 Chk1 inhibitors have been disclosed, including aryl- and heteroaryl-substituted urea compounds described in U.S. Patent Application No. 10/087,715 and U.S. Provisional Patent Application Nos. 60/583,080 and 60/602,968; diaryl urea compounds described in U.S. Patent Publica-
10 tion No. 2004/0014765, U.S. Patent Publication No. 2003/199511, U.S. Patent Publication No. 2004/0014765, and WO 03/101444; methylxanthines and related compounds described in Fan et al., *Cancer Res.* 55:1649-54, 1995; ureidothiophenes described in WO 03/029241 and
15 WO 03/028731; N-pyrrolopyridinyl carboxamides described in WO 03/028724; antisense Chk1 oligonucleotides described in WO 01/57206 and U.S. Patent No. 6,211,164; Chk1 receptor antagonists described in WO 00/16781; heteroaromatic carboxamide derivatives described in
20 WO 03/037886; aminothiophenes described in WO 03/029242; (indazolyl)benzimidazoles described in WO 03/004488; benzimidazole quinolinones described in U.S. Patent Publication No. 2004/0092535 and WO 04/018419; hetero-
cyclic-hydroxyimino-fluorenes described in WO 02/16326;
25 scytoneman derivatives, such as scytonemin, described in U.S. Patent No. 6,495,586; heteroarylbenzamides described in WO 01/53274; indazoles described in WO 01/53268; indolacarbazoles described in Tenzer et al., *supra*; chromane derivatives described in WO 02/070515; paullones
30 described in Schultz et al., *J. Med. Chem.*, Vol:2909-19, 1999; indenopyrazoles described in WO 99/17769; flavones described in Sedlacek et al., *Int J. Oncol.*, 9:1143-68, 1996; peptide derivatives of peptide loop of serine threonine kinases described in WO 98/53050; oxindoles

described in WO 03/051838; diazepinoindolones described
in WO 04/063198; pyrimidines described in WO 04/048343;
urea compounds described in WO 04/014876; and pyrrolo-
carbazoles, benzofuroisindoles, and azacyclopenta-
5 fluorenes described in WO 03/091255.


However, a need still exists in the art for
effective and selective inhibitors of Chk1. The present
invention addresses this and other needs.

SUMMARY OF THE INVENTION

10 The present invention relates to inhibitors of
the checkpoint kinase Chk1. The present Chk1 inhibitors
are useful in treating indications involving aberrant
cell proliferation, and as chemosensitizing and radiosens-
sitizing agents in the treatment of indications related
15 to DNA damage or lesions in DNA replication.

Therefore, one aspect of the present invention
is to provide compounds of structural formula (I). The
compounds are useful in a method of inhibiting Chk1 com-
prising a step of administering an effective amount of a
20 compound of structural formula (I) to an individual in
need thereof.

Compounds of formula (I) have a structural
formula:



(I),

25 wherein X¹ is null, -O-, -S-, -CH₂-, or
-N(R¹)-;

X² is -O-, -S-, or -N(R¹)-;

Y is O or S; or =Y represents two hydrogen atoms attached to a common carbon atom;

W is selected from the group consisting of heteroaryl, aryl, heterocycloalkyl, cycloalkyl, and
5 C₁₋₆alkyl substituted with a heteroaryl or aryl group, wherein said aryl group W is optionally substituted with one to four substituents represented by R², said heteroaryl group W is optionally substituted with one to four substituents represented by R⁵, and said heterocycloalkyl
10 and cycloalkyl groups W are optionally substituted with one or two C₁₋₆alkyl substituents;

R¹ is selected from the group consisting of hydro, C₁₋₆alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, and aryl;


R² is selected from the group consisting of
15 halo, optionally substituted C₁₋₆alkyl, C₂₋₆alkenyl, OCF₃, NO₂, CN, NC, N(R³)₂, OR³, CO₂R³, C(O)N(R³)₂, C(O)R³, N(R¹)COR³, N(R¹)C(O)OR³, N(R¹)C(O)C₁₋₆alkyleneC(O)R³, N(R¹)C(O)C₁₋₆alkyleneC(O)OR³, N(R¹)C(O)C₁₋₆alkyleneOR³, N(R¹)C(O)C₁₋₆alkyleneNHC(O)OR³, N(R¹)C(O)C₁₋₆alkyleneSO₂NR³,
20 C₁₋₆alkyleneOR³, and SR³;

R³ is selected from the group consisting of hydro, halo, C₁₋₆alkyl, C₂₋₆alkenyl, cycloalkyl, aryl, heteroaryl, SO₂R⁴, C₁₋₆alkyl substituted with one or more of halo, hydroxy, aryl, heteroaryl, heterocycloalkyl,
25 N(R⁴)₂, and SO₂R⁴, C₁₋₆alkylenearyl, C₁₋₆alkyleneheteroaryl, C₁₋₆alkyleneC₃₋₈heterocycloalkyl, C₁₋₆alkyleneSO₂aryl, optionally substituted C₁₋₆alkyleneN(R⁴)₂, OCF₃, C₁₋₆alkyleneN(R⁴)₃⁺, C₃₋₈heterocycloalkyl, and CH(C₁₋₆alkylene-N(R⁴)₂)₂, or two R³ groups are taken together to form an
30 optionally substituted 3- to 8-membered aliphatic ring;

R⁴ is selected from the group consisting of hydro, C₁₋₆alkyl, cycloalkyl, aryl, heteroaryl, C₁₋₆alkylenearyl, and SO₂C₁₋₆alkyl, or two R⁴ groups are taken

together to form an optionally substituted 3- to 8-membered ring;

R^5 is selected from the group consisting of C_{1-6} alkyl, C_{2-6} alkynyl, aryl, heterocycloalkyl, $N(R^3)_2$, OR^3 ,
 5 halo, N_3 , CN, C_{1-6} alkylenearyl, C_{1-6} alkylene $N(R^3)_2$, $C(O)R^3$,
 $C(O)OR^3$, $C(O)N(R^3)_2$, $N(R^1)C(O)R^3$, $N(R^1)C(O)OR^3$, and



R^6 is $-C\equiv C-R^7$ or heteroaryl;

R^7 is selected from the group consisting of
 10 hydro, C_{1-6} alkyl, aryl, C_{1-6} alkylenearyl, heteroaryl,
 C_{1-6} alkyleneheteroaryl, alkoxy;

R^8 , R^9 , and R^{10} , independently, are selected
 from the group consisting of halo, optionally substituted
 C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, OCF_3 , CF_3 , NO_2 , CN, NC,
 15 $N(R^3)_2$, OR^3 , CO_2R^3 , $C(O)N(R^3)_2$, $C(O)R^3$, $N(R^1)COR^3$, $N(R^1)C(O)-$
 OR^3 , $N(R^8)C(O)OR^3$, $N(R^1)C(O)C_{1-3}$ alkylene $C(O)R^3$, $N(R^1)C(O)-$
 C_{1-3} alkylene $C(O)OR^3$, $N(R^1)C(O)C_{1-3}$ alkylene OR^3 , $N(R^1)C(O)-$
 C_{1-3} alkylene $NHC(O)OR^3$, $N(R^1)C(O)C_{1-3}$ alkylene SO_2NR^3 , C_{1-3} alkyl-
 ene OR^3 , and SR^3 ;

20 or a pharmaceutically acceptable salt, or
 prodrug, or solvate thereof.

Another aspect of the present invention is to
 provide pharmaceutical compositions comprising one or
 more compound of structural formula (I), and use of the
 25 compositions in a therapeutic treatment of an indication,
 wherein inhibition of Chk1, in vivo or ex vivo, provides
 a therapeutic benefit or is of research or diagnostic
 interest.

Yet another aspect of the present invention is to provide a method of sensitizing cells in a subject undergoing a chemotherapeutic or radiotherapeutic treatment for an indication comprising administration of a compound of structural formula (I) in combination with a chemotherapeutic agent, a radiotherapeutic agent, or both, to the individual. A nonlimiting indication treated by this method is a cancer.

Another aspect of the present invention is to provide a method of inhibiting or preventing aberrant cell proliferation. In one embodiment, a method comprises contacting a cell population comprising aberrantly proliferating cells with at least one Chk1 activator in an amount and for a time sufficient to substantially synchronize cell cycle arrest among the aberrantly proliferating cells. Upon achieving substantial synchronization of cell cycle arrest in the cell population, the cell population is contacted with at least one Chk1 inhibitor in an amount and for a time sufficient to substantially abrogate the cell cycle arrest.

Another aspect of the present invention is to provide an article of manufacture for human pharmaceutical use comprising:

- (a) a pharmaceutical composition comprising a compound of structural formula (I);
- (b) a package insert informing that the composition is useful in the treatment of indications involving aberrant cell proliferation; and, optionally,
- (c) a container.

Another aspect of the present invention is to provide:

- (a) pharmaceutical composition comprising a compound of structural formula (I);


(b) a package insert informing that the composition is useful as a chemosensitizer or radiosensitizer in a treatment of an indication related to DNA lesions or DNA replication; and, optionally,

5 (c) a container.

These and other aspects of the present invention will become apparent from the following detailed description.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

10 Compounds of the present invention have a structural formula (I):



(I),

wherein X¹ is null, -O-, -S-, -CH₂-, or -N(R¹)-;

15 X² is -O-, -S-, or -N(R¹)-;

Y is O or S; or =Y represents two hydrogen atoms attached to a common carbon atom;

W is selected from the group consisting of heteroaryl, aryl, heterocycloalkyl, cycloalkyl, and C₁₋₆alkyl substituted with a heteroaryl or aryl group, wherein said aryl group W is optionally substituted with one to four substituents represented by R², said heteroaryl group W is optionally substituted with one to four substituents represented by R⁵, and said heterocycloalkyl and cycloalkyl groups W are optionally substituted with one or two C₁₋₆alkyl substituents;

20
25


R^1 is selected from the group consisting of hydro, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, and aryl;

R^2 is selected from the group consisting of halo, optionally substituted C_{1-6} alkyl, C_{2-6} alkenyl, OCF_3 ,
 5 NO_2 , CN , NC , $N(R^3)_2$, OR^3 , CO_2R^3 , $C(O)N(R^3)_2$, $C(O)R^3$,
 $N(R^1)COR^3$, $N(R^1)C(O)OR^3$, $N(R^1)C(O)C_{1-6}alkyleneC(O)R^3$,
 $N(R^1)C(O)C_{1-6}alkyleneC(O)OR^3$, $N(R^1)C(O)C_{1-6}alkyleneOR^3$,
 $N(R^1)C(O)C_{1-6}alkyleneNHC(O)OR^3$, $N(R^1)C(O)C_{1-6}alkyleneSO_2NR^3$,
 $C_{1-6}alkyleneOR^3$, and SR^3 ;

10 R^3 is selected from the group consisting of hydro, halo, C_{1-6} alkyl, C_{2-6} alkenyl, cycloalkyl, aryl, heteroaryl, SO_2R^4 , C_{1-6} alkyl substituted with one or more of halo, hydroxy, aryl, heteroaryl, heterocycloalkyl, $N(R^4)_2$, and SO_2R^4 , $C_{1-6}alkylenearyl$, $C_{1-6}alkyleneheteroaryl$,
 15 $C_{1-6}alkyleneC_{3-8}heterocycloalkyl$, $C_{1-6}alkyleneSO_2aryl$, optionally substituted $C_{1-6}alkyleneN(R^4)_2$, OCF_3 , $C_{1-6}alkyleneN(R^4)_3$, $C_{3-8}heterocycloalkyl$, and $CH(C_{1-6}alkyleneN(R^4)_2)_2$, or two R^3 groups are taken together to form an optionally substituted 3- to 8-membered aliphatic ring;

20 R^4 is selected from the group consisting of hydro, C_{1-6} alkyl, cycloalkyl, aryl, heteroaryl, $C_{1-6}alkylenearyl$, and $SO_2C_{1-6}alkyl$, or two R^4 groups are taken together to form an optionally substituted 3- to 8-membered ring;

25 R^5 is selected from the group consisting of C_{1-6} alkyl, $C_{2-6}alkynyl$, aryl, heterocycloalkyl, $N(R^3)_2$, OR^3 , halo, N_3 , CN , $C_{1-6}alkylenearyl$, $C_{1-6}alkyleneN(R^3)_2$, $C(O)R^3$, $C(O)OR^3$, $C(O)N(R^3)_2$, $N(R^1)C(O)R^3$, $N(R^1)C(O)OR^3$, and



R^6 is $-C\equiv C-R^7$ or heteroaryl;

R^7 is selected from the group consisting of hydro, C_{1-6} alkyl, aryl, C_{1-6} alkylenearyl, heteroaryl, C_{1-6} alkyleneheteroaryl, alkoxy;

5 R^8 , R^9 , and R^{10} , independently, are selected from the group consisting of halo, optionally substituted C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, OCF_3 , CF_3 , NO_2 , CN , NC , $N(R^3)_2$, OR^3 , CO_2R^3 , $C(O)N(R^3)_2$, $C(O)R^3$, $N(R^1)COR^3$, $N(R^1)C(O)-OR^3$, $N(R^8)C(O)OR^3$, $N(R^1)C(O)C_{1-3}$ alkylene $C(O)R^3$, $N(R^1)C(O)-C_{1-3}$ alkylene $C(O)OR^3$, $N(R^1)C(O)C_{1-3}$ alkylene OR^3 , $N(R^1)C(O)-C_{1-3}$ alkylene $NHC(O)OR^3$, $N(R^1)C(O)C_{1-3}$ alkylene SO_2NR^3 , C_{1-3} alkylene OR^3 , and SR^3 ;

and a pharmaceutically acceptable salt, or prodrug, or solvate thereof.


15 Preferred compounds of the present invention are those wherein X^1 and X^2 are $-N(H)-$;

Y is O or S ; and

W is optionally substituted heteroaryl. In one embodiment, W is heteroaryl containing at least two
20 heteroatoms selected from the group consisting of N , O , and S , said heteroaryl ring optionally substituted with one to four substituents selected from the group consisting of optionally substituted C_{1-6} alkyl, aryl, $N(R^3)_2$, OR^3 , $C(O)N(R^3)_2$, CO_2R^3 , CN , and halo, wherein R^3 is as
25 previously defined.


Other preferred compounds of structural formula (I) are those wherein W is selected from the group consisting of pyridazinyl, pyrimidinyl, pyrazinyl, and triazinyl, optionally substituted with one to four
30 substituents selected from the group consisting of C_{1-6} alkyl, aryl, $N(R^3)_2$, $C(O)N(R^3)_2$, CO_2R^3 , OR^3 , and halo.

In some preferred embodiments, W is selected from the group consisting of




optionally substituted with one to four substituents selected from the group consisting of C_{1-6} alkyl, C_{2-6} alkynyl, aryl, heteroaryl, CN, CO_2R^3 , $N(R^3)_2$, OR^3 , and halo.


In more preferred embodiments, W is



In a most preferred embodiment, W is pyrazinyl and X^1 and X^2 each are N(H).

In yet another preferred embodiment, R^6 is heteroaryl selected from the group consisting of





and optionally substituted with C₁₋₃alkyl-

$$5 \quad \text{eneN}(\mathbb{R}^4)_2.$$

As used herein, the term "alkyl" means straight chained and branched hydrocarbon groups containing the indicated number of carbon atoms, typically methyl, ethyl, and straight chain and branched propyl and butyl groups. Unless otherwise indicated, the hydrocarbon group can contain up to 20 carbon atoms. The term "alkyl" includes "bridged alkyl," i.e., a C₆-C₁₆ bicyclic or polycyclic hydrocarbon group, for example, norbornyl, adamantyl, bicyclo[2.2.2]octyl, bicyclo[2.2.1]heptyl, bicyclo[3.2.1]octyl, or decahydronaphthyl. Alkyl groups optionally can be substituted, for example, with hydroxy (OH), halo, aryl, heteroaryl, cycloalkyl, heterocycloalkyl, amino (N(R³)₂), and sulfonyl (SO₂R³), wherein R³ is as previously defined.

20 The term "cycloalkyl" means a cyclic C₃₋₈hydro-
carbon group, e.g., cyclopropyl, cyclobutyl, cyclohexyl,
or cyclopentyl. "Heterocycloalkyl" is defined similarly
as cycloalkyl, except the ring contains one to three
heteroatoms independently selected from the group con-
25 sisting of oxygen, nitrogen, and sulfur. Cycloalkyl and
heterocycloalkyl groups can be saturated or partially

unsaturated ring systems optionally substituted with, for example, one to three groups, independently selected from the group consisting of C₁₋₆alkyl, C₁₋₃alkyleneOH, C(O)NH₂, NH₂, oxo (=O), aryl, trifluoroethanoyl, and OH. Hetero-
5 cycloalkyl groups optionally can be further N-substituted with C₁₋₆alkyl, hydroxyC₁₋₆alkyl, C₁₋₃alkylenearyl, or C₁₋₃alkyleneheteroaryl.

The term "alkenyl" defined identically as "alkyl," except the group contains a carbon-carbon double
10 bond.

The term "alkynyl" is defined identically as "alkyl," except the group contains a carbon-carbon triple bond.

The term "alkylene" means an alkyl group having a substituent. For example, the term "C₁₋₆alkylene-C(O)OR" refers to an alkyl group containing one to six carbon atoms substituted with a -C(O)OR group. The alkylene group is optionally substituted with one or more substituent previously listed as an optional alkyl sub-
15 stituent.
20

The term "halo" or "halogen" means fluorine, bromine, chlorine, and iodine.

The term "aryl," alone or in combination, means a monocyclic or polycyclic aromatic group, preferably a monocyclic or bicyclic aromatic group, e.g., phenyl or naphthyl. Unless otherwise indicated, an aryl group can be unsubstituted or substituted with one or more, and in particular one to four groups independently selected from, for example, halo, C₁₋₆alkyl, C₂₋₆alkenyl,
25 OCF₃, NO₂, CN, NC, N(R³)₂, OR³, CO₂R³, C(O)N(R³)₂, C(O)R³, N(R¹)COR³, N(R¹)C(O)OR³, N(R¹)C(O)OR³, N(R¹)C(O)C₁₋₃alkyleneC(O)R³, N(R¹)C(O)C₁₋₃alkyleneC(O)OR³, N(R¹)C(O)C₁₋₃alkyleneOR³, N(R¹)C(O)C₁₋₃alkyleneNHC(O)OR³, N(R¹)C(O)-
30 C₁₋₃alkyleneSO₂NR³, C₁₋₃alkyleneOR¹, and SR³, wherein R¹ and

R³ are as previously defined. Exemplary aryl groups include, but are not limited to, phenyl, naphthyl, tetrahydronaphthyl, chlorophenyl, methylphenyl, methoxyphenyl, trifluoromethylphenyl, nitrophenyl, 2,4-methoxychloro-
5 phenyl, and the like. The terms "arylC₁₋₃alkyl" and "heteroarylC₁₋₃alkyl" mean an aryl or heteroaryl group having a C₁₋₃alkyl substituent.

The term "heteroaryl" means a monocyclic or bicyclic ring system containing one or two aromatic rings
10 and containing at least one nitrogen, oxygen, or sulfur atom in an aromatic ring. Unless otherwise indicated, a heteroaryl group can be unsubstituted or substituted with one or more, and in particular one to four, substituents selected from, for example, C₁₋₆alkyl, aryl, heteroaryl,
15 CF₃, CN, C(O)N(R³)₂, CO₂R², N(R³)₂, OR³, and halo, wherein R³ is as previously defined. Examples of heteroaryl groups include, but are not limited to, thienyl, furyl, pyridyl, oxazolyl, quinolyl, isoquinolyl, indolyl, triazinyl, triazolyl, isothiazolyl, isoxazolyl, imid-
20 izolyl, benzothiazolyl, pyrazinyl, pyrimidinyl, thiazolyl, and thiadiazolyl.

The term "hydro" is -H.

The term "hydroxy" is -OH.

The term "nitro" is -NO₂.

25 The term "cyano" is -CN.

The term "isocyano" is -NC.

The term "trifluoromethoxy" is -OCF₃.


The term "azido" is -N₃.

The term "3- to 8-membered ring" means carbo-
30 cyclic and heterocyclic aliphatic or aromatic groups, including, but not limited to, morpholinyl, piperidinyl, phenyl, thiophenyl, furyl, pyrrolyl, imidazolyl, pyrimidinyl, and pyridinyl, optionally substituted with


one or more, and in particular one to three, groups exemplified above for aryl groups.

The carbon atom content of hydrocarbon-containing moieties is indicated by a subscript designating the minimum and maximum number of carbon atoms in the moiety, e.g., "C₁₋₆alkyl" refers to an alkyl group having one to six carbon atoms, inclusive.

In the structures herein, for a bond lacking a substituent, the substituent is methyl, for example,



When no substituent is indicated as attached to a carbon atom on a ring, it is understood that the carbon atom contains the appropriate number of hydrogen atoms. In addition, when no substituent is indicated as attached to a carbonyl group or a nitrogen atom, for example, the substituent is understood to be hydrogen, e.g.,



The abbreviation "Me" is methyl. The abbreviation CO and C(O) is carbonyl (C=O).

The notation N(R^x) (wherein x represents an alpha or numeric character, such as for example R^a, R^b, R³, R⁴, and the like) is used to denote two R^x groups attached to a common nitrogen atom. When used in such notation, the R^x group can be the same or different, and are selected from the group as defined by the R^x group.

DNA-damaging agents that activate cell cycle checkpoints generally are referred to herein as "check-

point activators." DNA-damaging agents that activate the checkpoint designated "Chk1" (pronounced "check-one") are referred to herein as "Chk1 activators." Likewise, inhibitors of such checkpoints are referred to herein as

5 "checkpoint inhibitors" and "Chk1 inhibitors," respectively.

As used herein, Chk1 inhibitors are compounds that are capable of at least partially abrogating cell cycle checkpoint activity of the Chk1 protein. Abroga-

10 tion of cell cycle checkpoint is achieved when the cellular checkpoint mechanism is overcome sufficiently to allow the cell to pass from the cell cycle phase in which it is halted to the next phase in the cell cycle or to allow the cell to pass directly to cell death. Abroga-

15 tion of the cell cycle checkpoint permits cells to carry damage or imperfections to subsequent cell cycle phases, thereby inducing or promoting cell death. Cell death can occur by any associated mechanism, including apoptosis and mitotic catastrophe. The compounds of the invention

20 are Chk1 inhibitors.

Chk1 activator includes any known or after-discovered agent having the ability to activate Chk1 kinase activity, and thus induce at least partial cell cycle arrest. Chk1 activators include agents capable of

25 arresting the cell cycle at any phase of the cell cycle, which phase may be referred to herein as the "target phase" for that activator. Target phases include any of the cell cycle phases except mitosis, i.e., the G1 phase, S phase, and G2 phase. Chk1 activators useful in the

30 invention include DNA damaging agents, such as chemotherapeutic agents and/or radiation. Radiation Chk1 activators include, but are not limited to, ionizing radiation. Ionizing radiation includes electromagnetic or particulate radiation capable of producing ion pairs by inter-

acting with matter. Ionizing radiation includes X and gamma rays, alpha and beta particles, neutrons, and charged nuclei. Radiation includes ultraviolet light, visible light, infrared radiation, microwave radiation, and mixtures thereof. Assays such as that described in Example 9 can be used to determine whether an agent is a Chk1 activator.

"Inhibiting aberrant cell proliferation" means retarding the rate at which aberrantly proliferating cells proliferate or eliminating such proliferation altogether. This inhibition can result either from a decreased rate of replication, an increased rate of cell death, or both. Cell death can occur by any mechanism, including apoptosis and mitotic catastrophe.

"Preventing aberrant cell proliferation" means inhibiting aberrant cell proliferation prior to occurrence, or inhibiting the recurrence thereof.

"In vivo" means within a living subject, as within an animal or human. In this context, agents can be used therapeutically in vivo to retard or eliminate the proliferation of aberrantly replicating cells. The agents also can be used in vivo as a prophylactic to prevent aberrant cell proliferation or the manifestation of symptoms associated therewith.

"Ex vivo" means outside a living subject. Examples of ex vivo cell populations include cell cultures and biological samples, such as fluid or tissue samples from humans or animals. Such samples can be obtained by methods well known in the art. Exemplary biological fluid samples include blood, cerebrospinal fluid, urine, and saliva. Exemplary tissue samples include tumors and biopsies thereof. In this context, the present compounds can be in numerous applications, both therapeutic and experimental.

"Radiosensitizer" means a compound administered to a human or other animal in a therapeutically effective amount to increase the sensitivity of cells to electromagnetic radiation and/or to promote the treatment of diseases treatable with electromagnetic radiation.

"Radiation" includes, but is not limited to, radiation having the wavelength of 10^{-20} to 100 meters.

The term "container" means any receptacle and closure therefor suitable for storing, shipping, dispensing, and/or handling a pharmaceutical product.

The term "package insert" means information accompanying the product that provides a description of how to administer the product, along with the safety and efficacy data required to allow the physician, pharmacist, and patient to make an informed decision regarding use of the product. The package insert generally is regarded as the "label" for a pharmaceutical product.

The present invention includes all possible stereoisomers and geometric isomers of the compounds of structural formula (I). The present invention includes not only racemic compounds, but optically active isomers as well. When a compound of structural formula (I) is desired as a single enantiomer, it can be obtained either by resolution of the final product or by stereospecific synthesis from either an isomerically pure starting material or use of a chiral auxiliary reagent, for example, see Ma et al., *Tetrahedron: Asymmetry*, 8(6), pages 883-88, 1997. Resolution of the final product, an intermediate, or a starting material can be achieved by any suitable method known in the art. Additionally, in situations where tautomers of the compounds of structural formula (I) are possible, the present invention is intended to include all tautomeric forms of the compounds. As demonstrated hereafter, specific stereoisomers can

exhibit an exceptional ability to inhibit Chk1 in combination with chemotherapeutic or radiotherapeutic treatments.

Prodrugs of compounds of structural formula (I) also can be used as the compound in a method of the present invention. It is well established that a prodrug approach, wherein a compound is derivatized into a form suitable for formulation and/or administration, then released as a drug *in vivo*, has been successfully employed to transiently (e.g., bioreversibly) alter the physico-chemical properties of the compound (see, H. Bundgaard, Ed., "Design of Prodrugs," Elsevier, Amsterdam, 1985; Silverman, "The Organic Chemistry of Drug Design and Drug Action," Academic Press, San Diego, chapter 8, 1992; Hillgren et al., *Med. Res. Rev.*, 15, 83, 1995).

Compounds of the present invention can contain one or more functional groups. The functional groups, if desired or necessary, can be modified to provide a pro-drug. Suitable prodrugs include, for example, acid derivatives, such as amides and esters. It also is appreciated by those skilled in the art that N-oxides can be used as prodrugs.

Compounds of the invention can exist as salts. Pharmaceutically acceptable salts of the compounds of the invention generally are preferred in the methods of the invention. As used herein, the term "pharmaceutically acceptable salts" refers to salts or zwitterionic forms of the compounds of structural formula (I). Salts of compounds of formula (I) can be prepared during the final isolation and purification of the compounds or separately by reacting the compound with an acid having a suitable cation. Suitable pharmaceutically acceptable cations include alkali metal (e.g., sodium or potassium) and alkaline earth metal (e.g., calcium or magnesium) cat-

ions. In addition, the pharmaceutically acceptable salts of compounds of structural formula (I) that contain a basic center are acid addition salts formed with pharmaceutically acceptable acids. Examples of acids which can be employed to form pharmaceutically acceptable salts include inorganic acids such as hydrochloric, hydrobromic, sulfuric, and phosphoric, and organic acids such as oxalic, maleic, succinic, malonic, and citric. Nonlimiting examples of salts of compounds of the invention include, but are not limited to, hydrochloride, hydrobromide, hydroiodide, sulfate, bisulfate, 2-hydroxyethanesulfonate, phosphate, hydrogen phosphate, acetate, adipate, alginate, aspartate, benzoate, butyrate, camphorate, camphorsulfonate, citrate, digluconate, glycerolphosphate, hemisulfate, heptanoate, hexanoate, formate, succinate, malonate, fumarate, maleate, methanesulfonate, mesitylenesulfonate, naphthylenesulfonate, nicotinate, oxalate, pamoate, pectinate, persulfate, 3-phenylpropionate, picrate, pivalate, propionate, trichloroacetate, trifluoroacetate, glutamate, bicarbonate, undecanoate, lactate, citrate, tartrate, gluconate, benzene sulphonate, and p-toluenesulphonate salts. In addition, available amino groups present in the compounds of the invention can be quaternized with methyl, ethyl, propyl, and butyl chlorides, bromides, and iodides; dimethyl, diethyl, dibutyl, and diamyl sulfates; decyl, lauryl, myristyl, and steryl chlorides, bromides, and iodides; and benzyl and phenethyl bromides. In light of the foregoing, any reference to compounds of the present invention appearing herein is intended to include compounds of structural formula (I) as well as pharmaceutically acceptable salts, solvates, or prodrugs thereof.

The compounds of the present invention can be therapeutically administered as the neat chemical, but it may be useful to administer the compounds as a pharmaceutical composition or formulation. Thus, the present invention provides a pharmaceutical composition comprising a compound of formula (I) together with a pharmaceutically acceptable diluent or carrier therefor. Also provided is a process of preparing a pharmaceutical composition comprising admixing a compound of formula (I) with a pharmaceutically acceptable diluent or carrier therefor.

Accordingly, the present invention further provides pharmaceutical formulations comprising a compound of structural formula (I), or a pharmaceutically acceptable salt, prodrug, or solvate thereof, together with one or more pharmaceutically acceptable carriers and, optionally, other therapeutic and/or prophylactic ingredients. The carriers are "acceptable" in the sense of being compatible with the other ingredients of the formulation and not deleterious to the recipient thereof. Such carriers may be found for example, in *Remington's Pharmaceutical Sciences*, 17th Ed., Mack Publishing Co., Easton, PA (1985).

Compounds of the invention exhibit good potency against Chk1. Potency typically is expressed as the concentration of a compound required to achieve a certain result. The greater the potency, the less compound required to perform its intended function. In vitro potency typically is expressed in terms of IC₅₀ values and measured using a dose-response assay. IC₅₀ values can be measured by contacting a sensitive assay system with a compound of interest over a range of concentrations, including concentrations at which no or minimal effect is observed, through higher concentrations at which partial effect is observed, to saturating con-

centrations at which a maximum effect is observed. Theoretically, such assays of the dose-response effect of inhibitor compounds can be described as a sigmoidal curve expressing a degree of inhibition as a function of concentration when plotted on a log scale. The curve also theoretically passes through a point at which the concentration is sufficient to reduce activity of the check-point enzyme to a level that is 50% that of the difference between minimal and maximal enzyme activity observed in the assay. This concentration is defined as the Inhibitory Concentration at 50% inhibition or IC_{50} value.

IC_{50} values can be determined using conventional biochemical (acellular) assay techniques or cell-based assay techniques well known to those of ordinary skill in the art. An example of such an assay is provided in Example 1 below.

Preferably, IC_{50} values are obtained by performing the relevant assay at least twice, with the IC_{50} value expressed as the average (arithmetic mean, or "mean") of the individual values obtained. More preferably, the assay is repeated from 3 to 10 (or more) times, with the IC_{50} value expressed as the mean of the values obtained. Most preferably, the assay is performed a number of times sufficient to generate a statistically reliable mean IC_{50} value, using statistical methods known to those of ordinary skill in the art.

Compounds of the invention, when assayed as described in Example 1 below, exhibit IC_{50} values of less than about 5 μ M, and down to about 0.1 nM. In some embodiments compounds demonstrate an IC_{50} value of about 550 nM or less, in other embodiments less than about 250 nM, in others less than about 200 nM, in others less than about 150 nM, in others less than about 100 nM, in others less than about 75 nM, in others less than about 50 nM,

and in others less than about 25 nM. In preferred embodiment, compounds of the invention exhibit selectivity for inhibiting Chk1 over other protein kinases. Selectivity may be advantageous in reducing adverse side effects and/or increasing therapeutic index.

"Selectivity" is expressed herein as "fold selectivity." In general, fold selectivity, as used herein, is the IC_{50} of a test compound for a comparison enzyme divided by the IC_{50} of a comparator enzyme. In particular, fold selectivity for a Chk1 inhibitor, as used herein, is the IC_{50} of a Chk1 inhibitor (a test compound) for Chk1 (the comparison enzyme) divided by the IC_{50} for a comparator enzyme. Comparator enzymes against which compounds of the invention may be measured include at least the following protein kinases: Cdc2, Chk2, CTAK, EphA1, EphA2, Erk1, FGFR1, FGFR4, IR, JNK1, c-Kit, p38alpha, p38beta, p38delta, Ros, Rse, Rsk2, TrkA, TrkB, protein kinase A, protein kinase C, pp60v-src, protein kinase B/Akt-1, p38MapK, p70S6K, calcium calmodulin-dependent kinase II, and abl tyrosine kinase. Assays for determining IC_{50} values for a test compound against a comparator enzyme are described in Example 2 and are well known to those of ordinary skill in the art. Preferred compounds of the invention exhibit at least about 20-fold selectivity over the aforementioned protein kinases tested.

Compounds and pharmaceutical compositions suitable for use in the present invention include those wherein the active ingredient is administered in an effective amount to achieve its intended purpose. More specifically, a "therapeutically effective amount" means an amount sufficient to treat an individual suffering an indication, or to alleviate the existing symptoms of the indication. Determination of a therapeutically effective

amount is well within the capability of those skilled in the art, especially in light of the detailed disclosure provided herein.

In addition to the Chk1 inhibitor, pharmaceutical compositions of the invention can be formulated to include cytokines, lymphokines, growth factors, other hematopoietic factors, or mixtures thereof, to reduce adverse side effects that can arise from, or be associated with, administration of the pharmaceutical composition alone. Alternatively, such biologically active agents may be included in a pharmaceutical composition of the invention to promote a desired therapeutic effect. Adjuvant biologically active pharmaceutical compositions useful in pharmaceutical compositions of the invention include, but are not limited to, M-CSF, GM-CSF, TNF, IL-1, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-11, IL-12, IL-13, IL-14, IL-15, IL-16, IL-17, IL-18, IFN, TNF, G-CSF, Meg-CSF, GM-CSF, thrombopoietin, stem cell factor, erythropoietin, angiopoietins, including Ang-1, Ang-2, Ang-4, Ang-Y, and/or the human angiopoietin-like polypeptide, vascular endothelial growth factor (VEGF), angiogenin, bone morphogenic protein-1 (BMP-1), BMP-2, BMP-3, BMP-4, BMP-5, BMP-6, BMP-7, BMP-8, BMP-9, BMP-10, BMP-11, BMP-12, BMP-13, BMP-14, BMP-15, BMP receptor IA, BMP receptor IB, brain derived neurotrophic factor, ciliary neurotrophic factor, ciliary neurotrophic factor receptor cytokine-induced neutrophil chemotactic factor 1, cytokine-induced neutrophil chemotactic factor 2, endothelial cell growth factor, endothelin 1, epidermal growth factor, epithelial-derived neutrophil attractant, fibroblast growth factor (FGF) 4, FGF 5, FGF 6, FGF 7, FGF 8, FGF 8b, FGF 8c, FGF 9, FGF 10, FGF acidic, FGF basic, glial cell line-derived neurotrophic factor receptor 1, glial cell line-derived

neutrophilic factor receptor 2, growth related protein, growth related protein, growth related protein, growth related protein, heparin binding epidermal growth factor, hepatocyte growth factor, hepatocyte growth factor receptor, insulin-like growth factor I, insulin-like growth factor receptor, insulin-like growth factor II, insulin-like growth factor binding protein, keratinocyte growth factor, leukemia inhibitory factor, leukemia inhibitory factor receptor, nerve growth factor nerve growth factor receptor, neurotrophin-3, neurotrophin-4, placenta growth factor, placenta growth factor 2, platelet-derived endothelial cell growth factor, platelet derived growth factor, platelet derived growth factor A chain, platelet derived growth factor AA, platelet derived growth factor AB, platelet derived growth factor B chain, platelet derived growth factor BB, platelet derived growth factor receptor, platelet derived growth factor receptor, pre-B cell growth stimulating factor, stem cell factor, stem cell factor receptor, transforming growth factor (TGF), TGF, TGF 1, TGF 1.2, TGF 2, TGF 3, TGF 5, latent TGF 1, TGF, binding protein I, TGF binding protein II, TGF binding protein III, tumor necrosis factor receptor type I, tumor necrosis factor receptor type II, urokinase-type plasminogen activator receptor, vascular endothelial growth factor, and chimeric proteins and biologically or immunologically active fragments thereof.

The compounds of structural formula (I) also can be conjugated or linked to auxiliary moieties that promote a beneficial property of the compound in a method of therapeutic use. Such conjugates can enhance delivery of the compounds to a particular anatomical site or region of interest (e.g., a tumor), enable sustained therapeutic concentrations of the compounds in target cells, alter pharmacokinetic and pharmacodynamic proper-

ties of the compounds, and/or improve the therapeutic index or safety profile of the compounds. Suitable auxiliary moieties include, for example, amino acids, oligopeptides, or polypeptides, e.g., antibodies such as
5 monoclonal antibodies and other engineered antibodies; and natural or synthetic ligands to receptors in target cells or tissues. Other suitable auxiliaries include fatty acid or lipid moieties that promote biodistribution and/or uptake of the compound by target cells (see, e.g.,
10 Bradley et al., *Clin. Cancer Res.*, 7:3229, 2001).

Formulations of the present invention can be administered in a standard manner for the treatment of the indicated diseases, such as by oral, parenteral, transmucosal (e.g., sublingual or via buccal adminis-
15 tration), topical, transdermal, rectal, or inhalation (e.g., nasal or deep lung inhalation) administration. Parenteral administration includes, but is not limited to intravenous, intraarterial, intraperitoneal, subcutaneous, intramuscular, intrathecal, and intraarticular modes
20 of administration. Parenteral administration also can be accomplished using a high pressure technique, like POWDERJECT™ (Powderject Pharmaceuticals PLC, Oxford, England).

For oral administration and buccal adminis-
25 tration, the composition can be in the form of tablets or lozenges formulated in conventional manner. For example, tablets and capsules for oral administration can contain conventional excipients such as binding agents (for example, syrup, acacia, gelatin, sorbitol, tragacanth,
30 mucilage of starch, or polyvinylpyrrolidone), fillers (for example, lactose, sugar, microcrystalline cellulose, maize-starch, calcium phosphate, or sorbitol), lubricants (for example, magnesium stearate, stearic acid, talc, polyethylene glycol or silica), disintegrants (for

example, potato starch or sodium starch glycolate), or wetting agents (for example, sodium lauryl sulfate). The tablets can be coated according to methods well known in the art.

5 Alternatively, compounds of the present invention can be incorporated into oral liquid preparations such as aqueous or oily suspensions, solutions, emulsions, syrups, or elixirs, for example. Moreover, formulations containing these compounds can be presented as a
10 dry product for constitution with water or other suitable vehicle before use. Such liquid preparations can contain conventional additives, for example suspending agents, such as sorbitol syrup, methylcellulose, glucose/sugar syrup, gelatin, hydroxyethylcellulose, hydroxypropyl-
15 methylcellulose, carboxymethylcellulose, aluminum stearate gel, and hydrogenated edible fats; emulsifying agents, such as lecithin, sorbitan monooleate, or acacia; nonaqueous vehicles (which can include edible oils), such as almond oil, fractionated coconut oil, oily esters,
20 propylene glycol, and ethyl alcohol; and preservatives, such as methyl or propyl p-hydroxybenzoate and sorbic acid.

 Such preparations also can be formulated as suppositories, e.g., containing conventional suppository
25 bases, such as cocoa butter or other glycerides. Compositions for inhalation typically can be provided in the form of a solution, suspension, or emulsion that can be administered as a dry powder or in the form of an aerosol using a conventional propellant, such as dichlorodi-
30 fluoromethane or trichlorofluoromethane. Topical and transdermal formulations comprise conventional aqueous or nonaqueous vehicles, such as eye drops, creams, ointments, lotions, and pastes, or are in the form of a medicated plaster, patch, or membrane.

Additionally, compositions of the present invention can be formulated for parenteral administration by injection or continuous infusion. Formulations for injection can be in the form of suspensions, solutions, or emulsions in oily or aqueous vehicles, and can contain formulation agents, such as suspending, stabilizing, and/or dispersing agents. Alternatively, the active ingredient can be in powder form for constitution with a suitable vehicle (e.g., sterile, pyrogen-free water) before use.

A composition of the present invention also can be formulated as a depot preparation. Such long acting formulations can be administered by implantation (for example, subcutaneously or intramuscularly) or by intramuscular injection. Accordingly, the compounds of the invention can be formulated with suitable polymeric or hydrophobic materials (e.g., an emulsion in an acceptable oil), ion exchange resins, or as sparingly soluble derivatives (e.g., a sparingly soluble salt).


For veterinary use, a compound of formula (I), or a pharmaceutically acceptable salt, prodrug, or solvent thereof, is administered as a suitably acceptable formulation in accordance with normal veterinary practice. The veterinarian can readily determine the dosing regimen and route of administration that is most appropriate for a particular animal. Animals treatable by the present compounds and methods include, but are not limited to, pets, livestock, show animals, and zoo specimens.

SYNTHETIC METHODS

Compounds of the present invention can be prepared by the following synthetic schemes. Starting materials can be obtained from commercial sources or prepared by well-established literature methods known to those of


ordinary skill in the art. The groups X, R¹, R³, R⁴, R⁵, R⁶, R⁷, R⁸, R⁹ and R¹⁰ are defined above unless otherwise noted below.

Scheme 1




As illustrated in Scheme 1, compounds of formula 4 can be prepared from compounds of formula 6 by treatment with a base, such as DIEA, and diphenyl phosphoryl azide. A typical solvent for this reaction is THF, and the reaction is performed behind a blast shield at room temperature over a one to twelve hour period.

Scheme 2




Scheme 2 shows an alternative synthesis of compounds of formula 5. Compounds of formula 3 are treated with compounds of formula 7, which is prepared according to Scheme 4. A useful, nonlimiting solvent is DMF, and the reaction temperature is maintained between room temperature and 60°C for about one to twelve hours.

Scheme 3




As demonstrated in Scheme 3, compounds of formula 7 can be prepared from compounds of formula 8 by treatment with an aryl chloroformate, such as phenyl chloroformate or p-nitrophenyl chloroformate, in the presence of a base, such as pyridine. Nonlimiting solvents used in this reaction include CH₂Cl₂ or pyridine, at temperatures from 0°C to room temperature.

Scheme 4

Scheme 4 shows an approach to compounds of formula 3. Compounds of formula 1 are converted to compounds of formula 2 by treatment with aryl boronic acids and a source of palladium(0) (for example, palladium tetrakis triphenylphosphine) in the presence of a basic aqueous solution, such as Na_2CO_3 , K_2CO_3 , or K_3PO_4 . Nonlimiting examples of solvents used in this reaction include THF, dioxane, or ethylene glycol dimethyl ether. The reaction typically is performed at temperatures between 0°C and 90°C for about 1 to 12 h. Compounds of formula 2 are converted to compounds of formula 3 in the presence of Pd/C, Pt/C, or zinc, for example. Examples of solvents used in this reaction include, but are not limited to, MeOH, EtOH, or HOAc. Alternatively, compounds of formula 1 can be used to arylate terminal alkynes, 11, using a catalyst, such as $\text{PdCl}_2(\text{PPH}_3)_2$, or any other source of palladium(0). Reactions typically are conducted at temperatures varying from room temperature to 90°C , in the presence of a base, such as triethylamine.


Furthermore, compounds of formula 1, where X is a triflate, i.e., tf, can be obtained from compounds of formula 9. Typical reagents include triflic anhydride or N-phenyl triflimide. The reaction typically is performed at temperatures between -10°C and room temperature. A nonlimiting example of a solvent is dichloromethane. Nonlimiting examples of bases are TEA or diisopropyl ethyl amine.

Scheme 6




10

9



10



5


Scheme 6 illustrates an alternative synthesis for compounds of formula 5. Compounds of formula 3 can be converted to compounds of formula 10 following procedures described in Scheme 2. Compounds of formula 10
5 then can be converted to compounds of formula 5 following procedures described in Scheme 4.

Nonlimiting examples of compounds of structural formula (I) are provided below, the synthesis of which were performed in accordance with the general procedures set forth below and in copending U.S. Patent Application
10 Publication No. 2003/0069284, incorporated herein by reference. Additional compounds of the invention can be prepared using the above general schemes, and the following specific syntheses, by a judicious solution of starting
15 materials.

Abbreviations used in the syntheses described herein are: hours (h), minutes (min), atmosphere (atm), deionized (DI), nitrogen (N_2), water (H_2O), magnesium sulfate ($MgSO_4$), hydrochloric acid (HCl), dimethyl sulfoxide (DMSO), diisopropyl azodicarboxylate (DIAD),
20 dichloro palladium bis triphenylphosphene ($PdCl_2(PPh_3)_2$), triethylamine (TEA), carbon dioxide (CO_2), methylene chloride (CH_2Cl_2), chloroform ($CHCl_3$), methanol (MeOH), ammonium hydroxide (NH_4OH), ammonium chloride (NH_4Cl),
25 deuterated chloroform ($CDCl_3$), tetrahydrofuran (THF), N-methylpyrrolidone (NMP), acetic acid (HOAc), sodium hydroxide (NaOH), ethyl acetate (EtOAc), ethanol (EtOH), dimethyl sulfoxide (DMSO), diethyl ether (Et_2O), p-toluene sulfonic acid (p-TsOH), sodium carbonate (Na_2CO_3),
30 sodium bicarbonate ($NaHCO_3$), nitric acid (HNO_3), sodium chloride (NaCl), saturated (sat'd), thin layer chromatography (TLC), potassium carbonate (K_2CO_3), potassium phosphate (K_3PO_4), palladium on carbon (Pd/C), potassium chloride (KCl), phosphate buffered saline (PBS), cuprous


iodide (Cu(I)I), sodium sulfate (Na₂SO₄), dimethylformamide (DMF), 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU), and N,N-diisopropylethylamine (DIEA).

Intermediate 1:



5-Methyl-pyrazine-2-carbonyl azide

To a stirred suspension of 5-methyl-pyrazine-2-carboxylic acid (25 g, 181 mmol) in 540 mL THF at room temperature under N₂ was added DIEA (31.7 mL, 181 mmol) resulting in a brown solution. Diphenyl phosphoryl azide (39.2 mL, 181 mmol) then was added dropwise as a solution in 50 mL THF over 1 h behind a blast shield. The reaction was allowed to stir overnight. The reaction then was rotary evaporated to a small volume at room temperature and partitioned between Et₂O (1 L) and H₂O (1 L). The H₂O layer was back extracted with 2 x 250 mL Et₂O, and the combined organics washed 2 x 1 L with sat'd Na₂CO₃. The organics were dried (MgSO₄), filtered, and concentrated to a solid mass, which was triturated with Et₂O to give the product as a yellow solid (15 g, 50%). Purer compound could be isolated by taking x g of the crude product in 20x mL of Et₂O, and treating with 1-2x g of decolorizing carbon at room temperature for a few minutes. After filtration and concentration, this material was homogeneous by TLC in EtOAc and pure white. The recovery was typically 65%.

Compound 1:

1-(5-Methyl-pyrazin-2-yl)-3-(5-methyl-2-pyridin-3-ylethynyl)-urea

5 **Step 1. tert-Butyl-dimethyl-(4-methyl-2-nitro-phenylethynyl)-silane**

To a stirred solution of 1-bromo-4-methyl-2-nitro-benzene (3.24 g, 15.0 mmol), $\text{PdCl}_2(\text{PPh}_3)_2$ (1.05 g, 1.5 mmol), and Cu(I)I (5.7 g, 30 mmol) was added TEA (40
10 mL) followed by TMS-acetylene (5.3 mL, 37.5 mmol). After stirring at 60°C for 12 h, the reaction was filtered, diluted with EtOAc (150 mL) and 10% Na_2CO_3 (150 mL). The organic layer was washed with brine, dried over MgSO_4 , filtered, and dried under reduced pressure to yield 4.0 g
15 of an amorphous black residue.

Step 2. 1-Ethynyl-4-methyl-2-nitro-benzene

To a stirred solution of tert-butyl-dimethyl-(4-methyl-2-nitro-phenylethynyl)-silane (3.5 g, 15 mmol) in MeOH (20 mL) was added NaOH (1.8 g, 45 mmol, in 5 mL
20 of H_2O). After stirring for 1 hour, the reaction was diluted with EtOAc (150 mL) and 10% Na_2CO_3 (150 mL). The layers were separated and the aqueous layer was extracted with EtOAc (1 by 75 mL). The organic layers were combined, dried over MgSO_4 , filtered, and dried under re-
25 duced pressure to yield 1.40 g (58% over two steps) of a brown solid.

Step 3. 3-(4-Methyl-2-nitro-phenylethynyl)-pyridine

To a stirred solution of 3-bromopyridine (608 mg, 3.85 mmol), $\text{PdCl}_2(\text{PPh}_3)_2$ (246 mg, 0.35 mmol), and Cu(I)I (733 mg, 3.85 mmol) was added TEA (10 mL) followed
5 by 1-ethynyl-4-methyl-2-nitro-benzene (564 mg, 3.5 mmol). After stirring at 60°C for 4 h, the reaction was filtered, and diluted with EtOAc (150 mL) and 10% Na_2CO_3 (150 mL). The organic layer was washed with brine, dried over MgSO_4 , filtered, and dried under reduced pressure. The
10 material was purified using a Biotage 40M cartridge eluting with hexanes/EtOAc (7/3) to yield a light brown oil.

Step 4. 5-Methyl-2-pyridin-3-ylethynyl-phenylamine


To a solution of 3-(4-methyl-2-nitro-phenyl-ethynyl)-pyridine (1 mmol) in MeOH (1 mL) was added 0.5
15 mL sat'd NH_4Cl followed by zinc dust (0.33 g, 5.0 mmol). The mixture was stirred for 10 min, then diluted with EtOAc (50 mL) and Na_2CO_3 (50 mL of 10% aqueous solution). The organic layer was dried over MgSO_4 , filtered, and
20 concentrated under reduced pressure to yield the desired material as a clear oil.

Step 5. 1-(5-Methyl-pyrazin-2-yl)-3-(5-methyl-2-pyridin-3-ylethynyl-phenyl)-urea

To a stirred solution of 5-methyl-pyrazine-2-carbonyl azide (compound 7) (163 mg, 1.0 mmol) in toluene
25 (4 mL) and previously heated to 90°C for 15 min, was added 5-methyl-2-pyridin-3-ylethynyl-phenylamine (1 mmol). The mixture was cooled to 65°C and stirred for 12 h. The reaction mixture then was cooled to room temper-
30 ature. Ethyl acetate was added to the organic layer, and washed with brine, dried over MgSO_4 . After filtration,

and evaporation under reduced pressure, an oil was isolated and purified on column chromatography (silica gel) eluted with EtOAc/hexanes 1:4. ¹H-NMR (400 MHz, CDCl₃) δ 11.39 (br s, 1H), 8.79 (s, 1H), 8.61 (d, 1H), 8.25 (s, 1H), 8.19 (s, 1H), 8.15 (s, 1H), 7.79 (d, 1H), 7.41 (m, 2H), 7.31 (m, 1H), 6.91 (d, 1H), 2.41 (s, 3H), 2.37 (s, 3H). LRMS (apci, positive) m/e 344.4 (M+1).

Compound 2



10 1-(5-Methyl-pyrazin-2-yl)-3-(5-methyl-2-pyridin-3-yl-phenyl)-urea

Step 1. 3-(4-Methyl-2-nitro-phenyl)-pyridine


4-Bromo-2-nitro-phenol (648 mg, 3.0 mmol) and 3-pyridyl boronic acid (387 mg, 3.15 mmol) were diluted with 5 mL of dioxane and placed under N₂. Potassium carbonate was diluted in 1 mL of H₂O and added to the reaction mixture. Tetrakis(triphenylphosphine) palladium (173 mg, 0.15 mmol) was added, then the reaction was heated to 70°C and stirred overnight. The reaction was allowed to cool to room temperature, then diluted with 30 mL of EtOAc and 30 mL of 10% Na₂CO₃. The organic layer was washed with brine, dried over MgSO₄, filtered, and concentrated under reduced pressure. The material was purified with a Biotage 40M cartridge eluting with hexanes/EtOAc, 1/1 to yield an off white solid.

Step 2. 5-Methyl-2-pyridin-3-yl-phenylamine

Prepared according to compound 1, Step 4 from
3-(4-methyl-2-nitro-phenyl)-pyridine.

**Step 3. 1-(5-Methyl-pyrazin-2-yl)-3-
5 (5-methyl-2-pyridin-3-yl-phenyl)-urea**

Prepared according to compound 1, Step 5 using
5-methyl-2-pyridin-3-yl-phenylamine and compound 7. ¹H-
NMR (400 MHz, CDCl₃) δ 11.00 (br s, 1H), 9.00 (s, 1H),
8.75 (s, 1H), 8.70 (d, 1H), 8.40 (s, 2H), 7.75 (d, 1H),
10 7.40 (m, 2H), 7.15 (d, 1H), 7.05 (d, 1H), 2.45 (s, 3H),
2.40 (s, 3H). LRMS (apci, positive) m/e 320.3 (M+1).

Compound 3

15 1-(5-Methyl-pyrazin-2-yl)-3-(5-methyl-2-pyridin-4-yl-
phenyl)-urea

Step 1. 4-(4-Methyl-2-nitro-phenyl)-pyridine

Prepared according to Compound 2, Step 1 using
4-pyridyl boronic acid and 4-bromo-2-nitro-phenol.


Step 2. 5-Methyl-2-pyridin-4-yl-phenylamine

20 Prepared according to compound 1, Step 4 using
4-(4-methyl-2-nitro-phenyl)-pyridine.

**Step 3. 1-(5-Methyl-pyrazin-2-yl)-3-
(5-methyl-2-pyridin-4-yl-phenyl)-urea**

Prepared according to compound 1, Step 5 using 5-methyl-2-pyridin-4-yl-phenylamine and compound 7. ¹H-NMR (400 MHz, CDCl₃) δ 10.90 (br s, 1H), 8.65 (d, 2H), 8.25 (s, 1H), 8.20 (s, 1H), 8.10 (br s, 1H), 7.45 (d, 2H), 7.33 (s, 1H), 7.15 (d, 1H), 7.05 (d, 1H), 2.48 (s, 3H), 2.46 (s, 3H). LRMS (apci, positive) m/e 320.0 (M+1).

10 Compound 4:




1-(5-Methyl-pyrazin-2-yl)-3-(2-oxazol-5-yl-phenyl)-urea

Step 1: 2-Oxazol-5-yl-phenylamine

2-Oxazol-5-yl-phenyl nitro (190 mgs, 1 mmol) was dissolved in 3 mL of EtOH at room temperature. A catalytic amount of Pearlman's catalyst was added and the hydrogenation was performed under 1 atm. After filtration over celite, the solution was evaporated under reduced pressure. A yellow solid was isolated.

**20 Step 2: 1-(5-Methyl-pyrazin-2-yl)-3-
(2-oxazol-5-yl-phenyl)-urea**

Prepared according to compound 1, Step 5 using 2-oxazol-5-yl-phenylamine and compound 7. LRMS (apci, positive) m/e 296.0 (M+1).

Compound 5:

1-(5-Methyl-pyrazin-2-yl)-3-(5-methyl-2-thiazol-2-yl-phenyl) urea

5 **Step 1: 4-Methyl-2-nitro-benzamide**

4-Methyl-2-nitro benzamide was obtained from 4-methyl-2-nitro benzoic acid by the procedure described in *J. Am.Chem. Soc*, 79:1389 (1957).

Step 2: 4-Methyl-2-nitro-thio benzamide

10 4-Methyl-2-nitro benzamide (180 mgs, 1 mmol) and Belleu's reagent (529 mgs, 1 mmol) were dissolved in 3 mL of THF under N₂. The suspension was stirred overnight. A yellow solution had formed. The solution was concentrated, redissolved in 20 mL of CH₂Cl₂, and fresh
15 grade silica gel was added. The solution was evaporated under reduced pressure, and the silica gel/absorbed compound were loaded into a Biotage ZIF unit. The compound was chromatographed on a Biotage 12M column with EtOAc:hexanes 3:7. The desired fractions were pooled and
20 concentrated to give the desired material as a dark yellow solid.

Step 3: 2-(4-Methyl-2-nitro-phenyl-thiazole)

4-Methyl-2-nitro-thio benzamide (37 mg, 0.18 mmol) was dissolved in HOAc (5 mL) with pTsoH (90 mg, 0.047 mmol) and 2-bromo-1,1-diethoxy ethane (48 mg, 0.24
25

mmol). The mixture was heated to 100°C for 1.5 hr. The desired product was isolated using the procedure described in *Chem. Pharm. Bull.*, 39(9):2323-2332 (1991).


Step 4: 2-(4-Methyl-2-amino-phenyl-thiazole)

5 2-(4-Methyl-2-nitro-phenyl-thiazole) (30 mgs, 0.13 mmol) was dissolved in EtOH (3 mL) with a catalytic amount of Pearlman's catalyst. The procedure described for compound 4, Step 1 was followed. The desired material was obtained in good yields.

10 **Step 5: 1-(5-Methyl-pyrazin-2-yl)-3-(5-methyl-2-thiazol-2-yl-phenyl) urea**

Prepared according to compound 1, Step 5 using 2-(4-methyl-2-amino-phenyl-thiazole) and compound 7. LRMS (apci, positive) m/e 326.0 (M+1).

15 **Compound 6:**



1-[2-(4-Dimethylaminomethyl-thiazol-2-yl)-5-methyl-phenyl]-3-(5-methyl-pyrazin-2-yl)-urea

Step 1: 2-(4-Methyl-2-nitro-phenyl-thiazole-4-carboxylic acid ethyl ester)

4-Methyl-2-nitro-thio benzamide (60 mgs, 0.30 mmol, prepared as described in compound 5, Step 2, was stirred in 1 mL absolute EtOH at room temperature under N₂. Ethyl bromo pyruvate (65 mgs, 0.33 mmol) was added,

and the resulting solution was heated to 70°C for 3 h. The reaction was diluted to 30 mL with EtOAc and washed with saturated NaHCO₃, and saturated NaCl. The organic layer was dried over MgSO₄, filtered and concentrated to
5 a yellow oil, which was used as is in the next step.

**Step 2: 2-(4-Methyl-2-nitro phenyl)thiazol-
4-yl methanol**

2-(4-Methyl-2-nitro-phenyl-thiazole-4-carbox-
ylic acid ethyl ester (292 mg, 1 mmol) was dissolved in 5
10 mL of absolute EtOH at room temperature in an open flask. Sodium borohydride (1 mmol, 38 mg) was added portionwise over several hours and the reaction was monitored by TLC (EtOAc:hexanes 2:3). After completion of the reaction, 2N HCl was added carefully with stirring. After 15 min,
15 the clear yellow solution was concentrated on the rota-vap, and the crude mixture was partitioned between EtOAc (60 mL) and water (60 mL). The organics were isolated and washed with sat NaHCO₃ and 60 mL of sat NaCl. The organics were dried over MgSO₄, filtered and concentrated
20 to provide an orange solid as the desired material.

**Step 3 2-(4-Methyl-2-nitro-phenyl)thiazole-
4-carbaldehyde**

2-(4-Methyl-2-nitro phenyl)thiazol-4-yl meth-
anol (297 mg, 1.18 mmol) was stirred in 5 mL of CH₂Cl₂ at
25 room temperature under N₂. The Dess-Martin reagent (500 mg, 1.18 mmol) was added as a solid. After 30 minutes, the reaction was complete. The reaction was diluted to 60 mL of CH₂Cl₂ and washed with 60 mL of 1N NaOH. The organics were dried over MgSO₄, filtered and concentrated
30 under reduced pressure. The crude material was purified on a biotage column 12M eluted with EtOAc/hexanes 1:4 to give the desired material.

Step 4: Dimethyl-[2-(4-methyl-2-nitro-phenyl)-thiazol-4-ylmethyl]amine

Dimethylamine (640 uL of a 2M solution in MeOH), sodium acetate, and sodium cyanoborohydride (56 mgs, 0.89 mmol) were stirred in 2.6 mL MeOH at room temperature under N₂. Glacial HOAc was added to adjust the pH to 7-8. 2-(4-Methyl-2-nitro-phenyl)thiazole-4-carbaldehyde (159 mgs, 0.64 mmol) then was added as a solution in 3.2 mL of MeOH. After 2 h, product formation was apparent by mass spectroscopy. The reaction was allowed to proceed overnight. At that time, acetone (500 uL) was added to quench any unreacted borohydride and the reaction acidified to pH<3. The reaction was concentrated. The residue was partitioned between Et₂O (30 mL) and H₂O (30 mL). The aqueous phase was extracted with Et₂O, then neutralized with 1N NaOH to pH 10. The aqueous phase was re-extracted with Et₂O. The combined organics were dried over MgSO₄, filtered and concentrated under reduced pressure to provide the desired material.

Step 5: Dimethyl-[2-(4-methyl-2-amino-phenyl)-thiazol-4-ylmethyl]amine

Prepared according to compound 5, Step 4 from dimethyl-[2-(4-methyl-2-nitro-phenyl)-thiazol-4-ylmethyl]amine.

Step 6: 1-[2-(4-Dimethylaminomethyl-thiazol-2-yl)-5-methyl-3-phenyl]-3-(5-methyl-pyrazin-2-yl)-urea

Prepared according to Compound 1, Step 5 from compound 7 and dimethyl-[2-(4-methyl-2-amino-phenyl)-thiazol-4-ylmethyl]amine. LRMS (apci, positive) m/e 383.0 (M+1).

THERAPEUTIC METHODS

Compounds of the present invention can be used to treat conditions involving aberrant cell proliferation. For example, the compounds can be used to potentiate the therapeutic effects of radiation and/or a chemotherapeutic agent used in the treatment of cancers and other cell proliferation indications involving eukaryotic cells, including those in humans and other animals. In general, the present compounds inhibit aberrantly proliferating cells, both cancerous and noncancerous. For example, compounds of the invention can be used to enhance treatment of tumors that are customarily treated with an antimetabolite, e.g., methotrexate or 5-fluorouracil (5-FU).

Use of compounds of the present invention can result in partial or complete regression of aberrantly proliferating cells, i.e., the reduction or elimination of such cells from the cell population. Thus, for example, when the population of aberrantly proliferating cells are tumor cells, compounds of the invention can be used to retard the rate of tumor growth, decrease the number of tumors, and/or induce partial or complete tumor regression.

Compounds of the present invention can be used *in vivo* or *ex vivo* when no aberrant cell proliferation has been identified or where no aberrant cell proliferation is ongoing, but when aberrant cell proliferation is suspected or expected. Compounds of the present invention also can be used when aberrant cell proliferation has been previously treated in order to prevent or inhibit recurrence of the same.

One method of the present invention comprises administration of a therapeutically effective amount of a present Chk1 inhibitor compound in combination with a

chemotherapeutic agent to an individual in need thereof. Alternatively, a method of the present invention comprises administration of a therapeutically effective amount of at least one of the present Chk1 inhibitors to
5 an individual in need thereof in combination with an antibody, e.g., herceptin, that has activity in inhibiting the proliferation of cancer cells.

Cancers, therefore, are susceptible to enhanced treatment by administration of a present Chk1
10 inhibitor in combination with a chemotherapeutic agent or an antibody. Cancers treatable by the present invention include carcinomas and sarcomas that are characterized by solid tumors, and cancers of the myeloid or lymphoid systems, including leukemias, lymphomas, and other cancers
15 that typically lack a tumor mass, but are distributed in the vascular or lymphoreticular systems. These cancers include, for example, colorectal cancers, head and neck cancers, pancreatic cancers, breast cancers, gastric cancers, bladder cancers, vulvar cancers, leukemias,
20 lymphomas, melanomas, renal cell carcinomas, ovarian cancers, brain cancers, osteosarcomas, and lung cancers.

Compounds of the present invention, therefore, are useful in cancers mediated by Chk1 activity. More particularly, Chk1 activity is associated with forms of
25 cancer including, but not limited to, adult and pediatric oncology, growth of solid tumors/malignancies, myxoid and round cell carcinoma, locally advanced tumors, metastatic cancer, human soft tissue sarcomas, including Ewing's sarcoma, cancer metastases, including lymphatic metas-
30 tases, squamous cell carcinoma, particularly of the head and neck, esophageal squamous cell carcinoma, oral carcinoma, blood cell malignancies, including multiple myeloma, leukemias, including acute lymphocytic leukemia, acute antilymphocytic leukemia, chronic lymphocytic leu-

kemia, chronic myelocytic leukemia, and hairy cell leukemia, effusion lymphomas (body cavity based lymphomas), thymic lymphoma lung cancer (including small cell carcinoma, cutaneous T cell lymphoma, Hodgkin's lymphoma, non-Hodgkin's lymphoma, cancer of the adrenal cortex, ACTH-producing tumors, non-small cell cancers, breast cancer, including small cell carcinoma and ductal carcinoma), gastrointestinal cancers (including stomach cancer, colon cancer, colorectal cancer, and polyps associated with colorectal neoplasia), pancreatic cancer, liver cancer, urological cancers (including bladder cancer, such as primary superficial bladder tumors, invasive transitional cell carcinoma of the bladder, and muscle-invasive bladder cancer), prostate cancer, malignancies of the female genital tract (including ovarian carcinoma, primary peritoneal epithelial neoplasms, cervical carcinoma, uterine endometrial cancers, vaginal cancer, cancer of the vulva, uterine cancer and solid tumors in the ovarian follicle), malignancies of the male genital tract (including testicular cancer and penile cancer), kidney cancer (including renal cell carcinoma, brain cancer (including intrinsic brain tumors, neuroblastoma, astrocytic brain tumors, gliomas, and metastatic tumor cell invasion in the central nervous system), bone cancers (including osteomas and osteosarcomas), skin cancers (including malignant melanoma, tumor progression of human skin keratinocytes, and squamous cell cancer), thyroid cancer, retinoblastoma, neuroblastoma, peritoneal effusion, malignant pleural effusion, mesothelioma, Wilms's tumors, gall bladder cancer, trophoblastic neoplasms, hemangiopericytoma, and Kaposi's sarcoma.

A compound of the present invention also can be used to radiosensitize cells. Diseases treatable with radiation include, but are not limited to neoplastic di-

seases, benign and malignant tumors, and cancerous cells. Radiation treatment employs electromagnetic radiation such as gamma-radiation (10^{-20} to 10^{-13} m), X-ray radiation (10^{-12} to 10^{-9} m), ultraviolet light (10 nm to 400 nm),
5 visible light (400 nm to 700 nm), infrared radiation (700 nm to 1.0 mm), and microwave radiation (1 mm to 30 cm).

Some cancer treatment protocols currently employ radiosensitizers activated by electromagnetic radiation, e.g., X-rays. Examples of X-ray-activated radiosensitizers include, but are not limited to, the following: metronidazole, misonidazole, desmethylnisonidazole, pimonidazole, etanidazole, nimorazole, mitomycin C, RSU 1069, SR 4233, EO9, RB 6145, nicotinamide, 5-bromodeoxyuridine (BUdR), 5-iododeoxyuridine (IUdR), bromideoxycytidine, fluorodeoxyuridine (FUdR), hydroxyurea, cisplatin, and therapeutically effective analogs and derivatives thereof.

Photodynamic therapy (PDT) of cancers employs visible light as the radiation activator of the sensitizing agent. Examples of photodynamic radiosensitizers include the following, but are not limited to: hematoporphyrin derivatives, PHOTOFRIN[®], benzoporphyrin derivatives, NPe6, tin etioporphyrin (SnET2), pheoborbid-a, bacteriochlorophyll-a, naphthalocyanines, phthalocyanines, zinc phthalocyanine, and therapeutically effective analogs and derivatives of the same.

Radiosensitizers can be administered in conjunction with a therapeutically effective amount of one or more compounds in addition to the Chk1 inhibitor, such
30 compounds including, but not limited to, compounds that promote the incorporation of radiosensitizers to the target cells, compounds that control the flow of therapeutics, nutrients, and/or oxygen to the target cells, chemotherapeutic agents that act on the tumor with or

without additional radiation, or other therapeutically effective compounds for treating cancer or other disease. Examples of additional therapeutic agents or methods that can be used in conjunction with radiosensitizers include, but are not limited to, 5-fluorouracil (5-FU), leucovorin, oxygen, carbogen, red cell transfusions, perfluorocarbons (e.g., FLUOSOLW[®]-DA), 2,3-DPG, BW12C, calcium channel blockers, pentoxifylline, antiangiogenesis compounds, hydralazine, and L-BSO.

Chemotherapeutic agents that can be used in combination with a compound of the present invention to treat a cancer include, but are not limited to, alkylating agents, antimetabolites, hormones and antagonists thereof, radioisotopes, antibodies, as well as natural products, and combinations thereof. For example, an inhibitor compound of the present invention can be administered with antibiotics, such as doxorubicin and other anthracycline analogs, nitrogen mustards, such as cyclophosphamide, pyrimidine analogs such as 5-fluorouracil, cisplatin, hydroxyurea, taxol and its natural and synthetic derivatives, and the like. As another example, in the case of mixed tumors, such as adenocarcinoma of the breast, where the tumors include gonadotropin-dependent and gonadotropin-independent cells, the compound can be administered in conjunction with leuprolide or goserelin (synthetic peptide analogs of LH-RH). Other antineoplastic protocols include the use of an inhibitor compound with another treatment modality, e.g., surgery or radiation, also referred to herein as "adjunct antineoplastic modalities." Additional chemotherapeutic agents useful in the invention include hormones and antagonists thereof, radioisotopes, antibodies, natural products, and combinations thereof. Examples of chemotherapeutic agents useful in methods

employing compounds of the present invention are listed in the following table.

TABLE 1	
<p>A) <u>Alkylating agents</u></p> <p>i) <u>Nitrogen mustards</u> mechlorethamine cyclophosphamide ifosfamide melphalan chloroambucil</p> <p>ii) <u>Nitrosoureas</u> carmustine (BCNU) lomustine (CCNU) semustine (methyl-CCNU)</p> <p>iii) <u>Ethylenimine/Methyl-melamine</u> triethylenemelamine (TEM) triethylene thiophosphoramide (thiotepa) hexamethylmelamine (HMM, altretamine)</p> <p>iv) <u>Alkyl sulfonates</u> busulfan</p> <p>v) <u>Triazines</u> dacarbazine (DTIC)</p>	<p>G) <u>Natural products</u></p> <p>i) <u>Antimitotic drugs</u></p> <p>ii) <u>Taxanes</u> paclitaxel vinca alkaloids vinblastine (VLB) vincristine vinorelbine Taxotere[®] (docetaxel) estramustine estramustine phosphate</p> <p>iii) <u>Epipodophylotoxins</u> etoposide teniposide</p> <p>iv) <u>Antibiotics</u> actinomycin D daunomycin (rubidomycin) doxorubicin (adriamycin) mitoxantroneidarubicin bleomycin splicamycin (mithramycin) mitomycin C. dactinomycin aphidicolin</p> <p>v) <u>Enzymes</u> L-asparaginase L-arginase</p>
<p>B) <u>Antimetabolites</u></p> <p>i) <u>Folic Acid analogs</u> methotrexate trimetrexate pemetrexed (multitargeted antifolate)</p> <p>ii) <u>Pyrimidine analogs</u> 5-fluorouracil fluorodeoxyuridine gemcitabine cytosine arabinoside (AraC, cytarabine) 5-azacytidine 2,2'-difluorodeoxy-cytidine</p> <p>iii) <u>Purine analogs</u> 6-mercaptopurine 6-thioguanine azathioprine 2'-deoxycoformycin (pentostatin) erythrohydroxynonyl-adenine (EHNA) fludarabine phosphate 2-chlorodeoxyadenosine (cladribine, 2-CdA)</p>	<p>H) <u>Radiosensitizers</u> metronidazole misonidazole desmethylmisonidazole pimonidazole etanidazole nimorazole RSU 1069 EO9 RB 6145 SR4233 nicotinamide 5-bromodeoxyuridine 5-iododeoxyuridine bromodeoxycytidine</p>

TABLE 1	
C) <u>Type I Topoisomerase inhibitors</u> camptothecin topotecan irinotecan	I) <u>Miscellaneous agents</u> i) <u>Platinum coordination complexes</u> cisplatin carboplatin oxaliplatin anthracenedione mitoxantrone ii) <u>Substituted urea</u> hydroxyurea iii) <u>Methylhydrazine derivatives</u> N-methylhydrazine (MIH) procarbazine iv) <u>Adrenocortical suppressant</u> mitotane (o,p'-DDD) ainoglutethimide
D) <u>Biological response modifiers</u> G-CSF GM-CSF	J) <u>Cytokines</u> interferon (α , β , γ) interleukin-2
E) <u>Differentiation agents</u> retinoic acid derivatives	K) <u>Photosensitizers</u> hematoporphyrin derivatives Photofrin benzoporphyrin derivatives Npe6 tin etioporphyrin (SnET2) pheoboride-a bacteriochlorophyll-a naphthalocyanines phthalocyanines zinc phthalocyanines
F) <u>Hormones and antagonists</u> i) <u>Adrenocorticosteroids/antagonists</u> prednisone and equivalents dexamethasone ainoglutethimide ii) <u>Progestins</u> hydroxyprogesterone caproate medroxyprogesterone acetate megestrol acetate iii) <u>Estrogens</u> diethylstilbestrol ethynyl estradiol and equivalents iv) <u>Anti-estrogen</u> tamoxifen v) <u>Androgens</u> testosterone propionate fluoxymesterone and equivalents vi) <u>Antiandrogens</u> flutamide gonadotropin-releasing hormone analogs leuprolide vii) <u>Nonsteroidal antiandrogens</u> flutamide	L) <u>Radiation</u> X-ray ultraviolet light gamma radiation visible light infrared radiation microwave radiation

Examples of chemotherapeutic agents that are particularly useful in conjunction with radiosensitizers include, for example, camptothecin, carboplatin, cis-platin, daunorubicin, doxorubicin, interferon (α , β , γ), irinotecan, hydroxyurea, chlorambucil, 5-fluorouracil, methotrexate, 2-chloroadenosine, fludarabine, azacytidine, gemcitabine, pemetrexed, interleukin 2,

irinotecan, docetaxel, paclitaxel, topotecan, and therapeutically effective analogs and derivatives of the same.

In accordance with the present invention, compounds of the present invention are useful in combination with gemcitabine, alone or further with paclitaxel. Compounds of the present invention also are useful in combination with pemetrexed, alone or further with cisplatin, carboplatin, or other platins. A present Chk1 inhibitor also can be administered in combination with gemcitabine and pemetrexed.

A present Chk1 inhibitor administered in combination with gemcitabine can be useful in the treatment of, for example, pancreatic carcinoma, leiomyosarcoma of the uterus, bone sarcoma, metastatic nonsmall cell lung cancer, extremity and trunk soft tissue sarcoma, renal cell cancer, adenocarcinoma, and Hodgkin's disease. A present Chk1 inhibitor administered with pemetrexed can be useful in the treatment of mesothelioma.

Compounds of the present invention also can potentiate the efficacy of drugs used in the treatment of inflammatory diseases, conditions, or disorders characterized by aberrant cell proliferation. Examples of inflammatory diseases that can be treated with compounds of the present invention include, but are not limited to, rheumatoid arthritis (RA), psoriasis, vitiligo, Wegener's granulomatosis, systemic-onset juvenile chronic arthritis (JCA), and systemic lupus erythematosus (SLE). Treatment of arthritis, Wegener's granulomatosis, and SLE often involves the use of immunosuppressive therapies, such as ionizing radiation, methotrexate, and cyclophosphamide. Such treatments typically induce, either directly or indirectly, DNA damage. Inhibition of Chk1 activity within the offending immune cells render the cells more sensitive to control by these standard treatments. Psoriasis

and vitiligo commonly are treated with ultraviolet radiation (UV) in combination with a psoralen. The compounds of the present invention enhance the killing effect of UV and a psoralen, and increase the therapeutic index of this treatment regimen. In general, compounds of the present invention potentiate control of inflammatory disease cells when used in combination with immunosuppressive drugs.

The compound of the present invention also can be used in methods of treating other noncancerous conditions characterized by aberrantly proliferating cells. Such conditions include, but are not limited to, atherosclerosis, restenosis, vasculitis, nephritis, retinopathy, renal disease, proliferative skin disorders, psoriasis, keloid scarring, actinic keratosis, Stevens-Johnson Syndrome, osteoporosis, hyperproliferative diseases of the eye including epithelial down growth, proliferative vitreoretinopathy (PVR), diabetic retinopathy, Hemangio-proliferative diseases, ichthyosis, and papillomas.

One preferred method of administering a Chk1 inhibitor of the present invention is described in WO 05/027907, the disclosure of which is incorporated by reference. Such methods for inhibiting aberrant cell proliferation involve scheduling administration of a Chk1 activator (e.g., a chemotherapeutic agent) and a Chk1 inhibitor according to the present invention. In this method, at least one Chk1 activator is administered at a dose and for a time sufficient to induce substantial synchronization of cell cycle arrest in proliferating cells. Upon achieving substantial phase synchronization, at least one Chk1 inhibitor is administered to abrogate the cell cycle arrest and induce therapeutic cell death. The method is useful with any Chk1 activator, and finds

application in treating or preventing cancerous and noncancerous conditions involving aberrant cell proliferation.

A population of aberrantly proliferating cells can be contacted with one, or more than one, Chk1 inhibitor of the invention. If more than one Chk1 inhibitor is used, the Chk1 inhibitors can be contacted with the cells using the same or different methods (e.g., simultaneously or sequentially, for the same or different durations, or by the same or different modalities) as determined by the skilled artisan, e.g., an attending physician (in the case of human patients) or a laboratory experimentalist (in the case of an *in vitro* or *ex vivo* procedure).

A population of aberrantly proliferating cells also can be contacted with one or more Chk1 activator. If more than one Chk1 activator is used, the Chk1 activators can be contacted with the cells using the same or different methods, generally as described in the context of Chk1 inhibitors above.

Compounds of the present invention can be applied to cell populations *ex vivo*. For example, the present compounds can be used *ex vivo* to obtain information concerning the optimal schedule and/or dosing for administering a Chk1 inhibitor for a given indication, cell type, patient, and/or other treatment parameter. This information can be used for experimental purposes or in a clinic to determine protocols for *in vivo* treatment. Other *ex vivo* uses for compounds of the present invention will be apparent to persons skilled in the art.

As appreciated by persons skilled in the art, additional active or ancillary agents may be used in the methods described herein. As also appreciated by persons skilled in the art, reference herein to treatment extends

to prophylaxis, as well as to treatment of established diseases or symptoms.

The amount of a compound of the invention required for use in treatment varies with the nature of the condition being treated, and with the age and the condition of the patient, and is ultimately determined by the attendant physician or veterinarian. In general, however, doses administered for adult human treatment typically are in the range of 0.001 mg/kg to about 100 mg/kg per day. The dose can be administered in a single dose, or as multiple doses administered at appropriate intervals, for example as two, three, four or more subdoses per day. In practice, the physician determines the dosing regimen suitable for an individual patient, and the dosage varies with the age, weight, and response of the particular patient. The above dosages are exemplary of the average case, but individual instances exist wherein higher or lower dosages are merited, and such are within the scope of the present invention.

Contact of the cell population with a present Chk1 inhibitor, at any dose, is for a time sufficient to achieve substantial abrogation of the cell cycle checkpoint. Typically, though not necessarily, such times include up to about 72 h to about 96 h, depending upon various factors. In some embodiments, it is desirable or necessary to administer Chk1 inhibitor over a period of up to about several weeks or more, as determined by the attending physician or technician. Thus, a present Chk1 inhibitor typically can be administered for up to about 1 hour, up to about 2 h, up to about 3 h, up to about 4 h, up to about 6 h, up to about 12 h, up to about 18 h, up to about 24 h, up to about 48 h, or up to about 72 h. Persons skilled in the art appreciate that the ranges of time expressed herein are merely exemplary and that

ranges and subranges within and outside those expressed also are within the scope of the invention.

Chk1 inhibitors of the present invention can be administered over a plurality of doses. For example, the Chk1 inhibitor can be given at a frequency of: four doses delivered as one dose per day at four-day intervals (q4d x 4); four doses delivered as one dose per day at three-day intervals (q3d x 4); one dose delivered per day at five-day intervals (qd x 5); one dose per week for three weeks (qwk3); five daily doses, with two days rest, and another five daily doses (5/2/5); or, any dose regimen determined to be appropriate for the circumstance.

EXAMPLES

Example 1

15 Determination of IC₅₀ Values of the Chk1 Inhibitors

Human Chk1 cDNA was identified and cloned as described previously in WO 99/11795, filed September 4, 1998. A FLAG[®] tag was inserted in frame with the amino terminus of the full-length Chk1. The 5' primer contains an EcoRI site, a Kozak sequence, and also encodes a FLAG[®] tag for affinity purification using the M2 Antibody (Sigma, Saint Louis, IL). The 3' primer contains a SalI site. The PCR-amplified fragment was cloned into pCI-Neo as an EcoRI-SalI fragment (Invitrogen, Carlsbad, CA), then subcloned as an EcoRI-NotI fragment into pFastBacI (Gibco-BRL, Bethesda, MD). Recombinant baculovirus was prepared as described in the Gibco-BRL Bac-to-Bac manual and used to infect Sf-9 cells grown in CCM3 medium (HyClone Laboratories, Logan, UT) for expression of FLAG[®]-tagged Chk1 protein.

FLAG[®]-tagged Chk1 was purified from frozen pellets of baculovirus-infected SF9 cells. Frozen cell pellets were mixed with an equal volume of 2X lysis buffer containing 100 mM Tris-HCl pH 7.5, 200 mM NaCl, 50 mM B-glycerophosphate, 25 mM NaF, 4 mM MgCl₂, 0.5 mM EGTA, 0.2% TWEEN[®]-20, 2 mM sodium vanadate, 2 mM DTT, and a cocktail of protease inhibitors (Complete mini, Boehringer Mannheim 2000 catalog #1836170). Cells then were dounced 20 times with the loose pestle of a dounce homogenizer and centrifuged at 48,400 x g for 1 hour. The M2 affinity was prewashed with 10 column volumes of 50 mM glycine pH 3.5 followed by 20 mM Tris pH 7.5, 150 mM NaCl alternating three times and ending with a Tris NaCl wash. The column then was washed with 25 column volumes of 20 mM Tris pH 7.5, 150 mM NaCl, 0.1% TWEEN[®]-20, 1 mM EGTA, 1 mM EDTA and 1X complete mini protease tablets. The cleared lysate then was bound to M2 affinity resin in batch at 4°C for 4 h. The mixture of resin and lysate then was poured into a column and the flow through collected. The resin was washed with 10 column volumes of 20 mM Tris pH 7.5, 150 mM NaCl, and 3 mM N-octyl glucoside. FLAG[®]-tagged Chk1 then was eluted from the column with 6 column volumes of cold 20 mM Tris pH 7.5, 150 mM NaCl, 3 mM N-octyl glucoside containing 0.5 mg/mL FLAG[®] peptide (Sigma, 2000 Catalog # F-3290). Three fractions were collected and analyzed for the presence of FLAG-tagged Chk1.

The assay for Chk1 kinase activity that includes 100 ng purified FLAG[®]-Chk1 (150 pmol of ATP/min), 20 μM Cdc25C peptide (H-leu-tyr-arg-ser-pro-ser-met-pro-glu-asn-leu-asn-arg-arg-arg-arg-OH) (SEQ ID NO: 1), 4 μM ATP, 2 μCi [³²P]γ-ATP, 20 mM Hepes pH 7.2, 5 mM MgCl₂, 0.1% NP40, and 1 mM DTT. Reactions were initiated by the addition of ATP-containing reaction mix and carried out

at room temperature for 10 min. Reactions were stopped by the addition of phosphoric acid (150 mM final concentration) and transferred to phosphocellulose discs. The phosphocellulose discs were washed five times with 150 mM
5 phosphoric acid and air-dried. Scintillation fluid was added and discs were counted in a Wallac scintillation counter. The assay was incubated in the presence of a broad range of concentrations of Chk1 inhibitor compound and an IC_{50} value for the compound was calculated. As
10 indicated above, all compounds of the invention subjected to the assay exhibited IC_{50} values in the assay of less than about 500 nM.

Example 2

Selectivity

15 Chk1 inhibitors of the present invention were tested for selectivity, with Chk1 as the comparison enzyme and the following protein kinases as comparator enzymes: Cdc2, Chk2, CTAK, EphA1, EphA2, Erk1, FGFR1, FGFR4, IR, JNK1, c-Kit, p38alpha, p38beta, p38delta, Ros,
20 Rse, Rsk2, TrkA, TrkB, protein kinase A, protein kinase C, pp60v-src, protein kinase B/Akt-1, p38MapK, p70S6K, calcium calmodulin-dependent kinase II, and abl tyrosine kinase.

The IC_{50} value of a compound versus Chk1 was
25 measured as described above. The IC_{50} value of the compound against comparator enzymes was measured using the SelectSmart™ (MDS Pharma Servies, Bothell, Washington, USA) proprietary technology platform with either a modified ELISA procedure or fluorescence polarization. All
30 inhibitors tested showed at least a 20-fold selectivity for Chk1 over the tested comparator enzymes.

Alternatively, assays for determining IC₅₀ for each of these kinases have been previously described in the literature, including U.S. Patent Publication No. 2002/016521 and WO 95/19988, both of which are incorporated by reference here.

Example 3

Chk1 Inhibitors of the Invention Inhibit Chk1 Function in Cells

To establish that the Chk1 inhibitors of the invention inhibit Chk1 function in cells, inhibitors can be tested in molecular cell-based assays. Because mammalian Chk1 has been shown to phosphorylate Cdc25C *in vitro*, suggesting that it negatively regulates cyclin B/cdc2 in response to DNA damage, the ability of the Chk1 inhibitors to enhance the activity of CyclinB/cdc2 can be analyzed. The experiment can be designed as follows: HeLa cells are irradiated with 800 rads and incubated for 7 h at 37°C. Because these cells are functionally p53 negative, they arrest exclusively in G2. Then, nocodazole is added to a concentration of 0.5 µg/mL and the cells are incubated for 15 h at 37°C. The addition of nocodazole is designed to trap any cells that progress through the G2 arrest into M. Finally, a Chk1 inhibitor is added for 8 h, the cells harvested, lysed and immunoprecipitated equal amounts of protein with an antibody to Cyclin B1 (New England Biolabs) as suggested by the manufacturer. Immunoprecipitates then are analyzed for Cyclin B-associated cdc2 kinase activity by assaying histone H1 kinase activity (Yu et al., *J Biol Chem.*, Dec. 11, 1998; 273(50):33455-64).

In addition, the ability of the subject Chk1 inhibitors to abrogate the ionizing radiation-induced G2

DNA damage checkpoint can be established using mitotic index assay experiments. HeLa cells (approximately 1×10^6) are treated as described above. Cells are harvested by centrifugation, washed once with PBS, then
5 resuspended in 2.5 mL of 75 mM KCl and centrifuged again. The cells then are fixed in 3 mL of freshly prepared cold HOAc:MeOH (1:3) and incubated on ice for 20 min. Cells are pelleted, fix solution aspirated and resuspended in 0.5 mL of PBS. Mitotic spreads are prepared by pipeting
10 100 μ L of the fixed cells onto a glass microscope slide and flooding the sample with 1 mL of fix solution. Slides then are air dried, stained with Wright's stain (Sigma) for 1 min, followed by one wash with H₂O and one wash with 50% MeOH. The presence of condensed chromo-
15 somes and lack of nuclear envelope identifies mitotic cells.

Example 4

Chk1 Inhibitors of the Present Invention Enhance Killing of Cells by Cancer Treatments

20 To demonstrate that the inhibition of Chk1 by a compound of the present invention sensitizes targeted cells to the killing effect of DNA-damaging agents, cells can be incubated in the presence of a present Chk1 inhibitor and exposed to either irradiation or a chemical DNA-
25 damaging agent. Cells plated at a density of 1000-2000 per well in 96-well microtitre plates are grown in RPMI 1640 containing 10% FBS, 100 U/mL penicillin and 100 μ g/mL streptomycin for 18 h at 37°C in a humidified incubator with 5% CO₂. Cells tested can include any cells
30 or cell lines of interest, such as HeLa, ACHN, 786-0, HCT116, SW620, HT29, Colo205, SK-MEL-5, SK-MEL-28, A549, H322, OVCAR-3, SK-OV-3, MDA-MB-231, MCF-7, PC-3, HL-60,

K562, and MOLT4. All cell line designations refer to the following human cell lines:

HeLa	cervical adenocarcinoma
ACHN	renal adenocarcinoma
786-0	renal adenocarcinoma
HCT116	colon carcinoma
SW620	colon carcinoma, lymph node metastasis
HT-29	colonrectal adenocarcinoma
Colo205	colon adenocarcinoma
SK-MEL-5	melanoma
SK-MEL-28	malignant melanoma
A549	lung carcinoma
H322	bronchoalveolar carcinoma
OVCAR-3	ovarian adenocarcinoma
SK-OV-3	ovarian adenocarcinoma
MDA-MB-231	breast adenocarcinoma
MCF-7	breast adenocarcinoma
PC-3	prostate adenocarcinoma, from metastasis to bone
HL-60	acute promyelocytic leukemia
K562	chronic myelogenous leukemia
MOLT4	acute lymphoblastic leukemia; T lymphoblast

5

Cells are treated with media containing chemotherapeutic drugs alone or chemotherapeutic drugs and a Chk1 inhibitor. Cells are incubated for approximately 5 days before growth is measured by determination of levels of ³H-thymidine uptake. Chemotherapeutic drugs include etoposide, doxorubicin, cisplatin, chlorambucil, 5-fluorouracil. The drug concentration necessary to inhibit cell growth to 90% of untreated control cells is defined as the GI₉₀.

15

Compounds of the present invention can be tested with additional antimetabolites, including methotrexate, hydroxyurea, 2-chloroadenosine, fludarabine, azacytidine, and gemcitabine to assess therein ability to enhance killing of the agents. Compounds of the present invention can be compared to one another by

20

assessing enhanced killing of HT29 colorectal carcinoma in combination with gemcitabine.

In addition, the ability of the Chk1 inhibitors of the invention to enhance killing by radiation can
5 be tested.

Example 5

Sensitive Assay to Measure Chk1 Inhibitor Activity in Animal Models

The following sensitive assay was developed to
10 measure Chk1 inhibitor activity in rodent tumor models.
In particular, the assay can be used, *inter alia*, to
measure the ability of a Chk1 inhibitor to block Chk1
function in the tumor model, and to allow for assessment
of conditions that facilitate access of the Chk1 inhibi-
15 tor to the molecular target.

The ability of selective Chk1 inhibitors to
abrogate a chemotherapy-induced checkpoint is measured
using a quantitative immunofluorescent assay that
measures mitotic index by monitoring histone H3 phos-
20 phorylation on serine 10 (H3-P), a mitosis-specific event
(Ajiro et al., *J Biol Chem.*, 271:13197-201, 1996; Goto et
al., *J Biol Chem.*, 274:25543-9, 1999). The assay proto-
col is as follows. Tumors from rodents treated or un-
treated with Chk1 activator (in the present study, chemo-
25 therapy agent) and/or Chk1 inhibitor, are excised and
paraffin embedded. The tumors are cut into 6 micron
thick slices and mounted on glass slides. The paraffin
is removed from the slides by 3 min successive treatments
with xylene, 100% ethanol, 95% EtOH, 70% EtOH, and DI
30 H₂O. The slides then are heated to 95°C in 10 mM sodium
citrate for 10 min followed by a 20 min cooling step.
The slides are blocked for 30 min with Block buffer (20%

normal human serum and 2% bovine serum albumin in phosphate buffered saline containing 0.05% Triton X-100 (PBST)). The antiphospho histone H3 antibody (Upstate Biotech, Cat. #06-570) is diluted 1:200 in the Block
5 buffer and incubated with the slides for 1 h. The slides are washed 3 times 5 min in PBST. The secondary antibody, donkey antirabbit rhodamine (Jackson, cat #711-295-152) is added for 30 min. The slides then are washed twice in PBST and 75 μ M of 0.1 μ M/ml DAPI (Sigma) in PBS
10 is added and allowed to stain for 30 min. The slides then are washed two more times in PBST and mounted with Vectashield (Vector, cat # H-1400). Slides are viewed using fluorescence microscopy. The percentage of cells stained with H3-P antibody relative to total (DAPI
15 stained) cells are quantified using Metamorph software (Universal Imaging Corporation, Version 4.6).

Example 6

Selective Chk1 Inhibitors Abrogate DNA Damage-Induced G2 and S Phase Checkpoints

20 Previous studies demonstrated that selective Chk1 inhibitors substantially abrogate the DNA damage-induced G2/M and S phase checkpoints. In the former, DNA damage is induced by ionizing radiation (IR), whose target phase is the G2 phase. In the latter, DNA damage is
25 induced by chemotherapeutic agents whose target phase is the S phase. See published U.S. Patent Application Publication 2003/0069284 and references cited therein.

Briefly, Chk1 inhibitor abrogation of IR-induced G2 DNA damage checkpoint is assayed by mitotic
30 index experiments. Approximately 1×10^6 HeLa cells are irradiated with 800 rads and incubated for 7 h at 37°C. Because these cells are functionally p53 negative, they

arrest exclusively in G2. Nocodazole then is added to a concentration of 0.5 $\mu\text{g/mL}$ and incubated for 15 h at 37° C. (The addition of nocodazole is designed to trap cells that progressed through the G2 arrest in mitosis thus preventing them from further progressing into G1 and allowing for quantification of M phase cells.) A selective Chk1 inhibitor is added for 8 h, and the cells are harvested by centrifugation, washed once with PBS, then resuspended in 2.5 mL 75 mM KCl and centrifuged again.

10 The cells then are fixed in 3 mL of freshly prepared cold HOAc:MeOH (1:3) and incubated on ice for 20 min. Cells are pelleted, the fix solution is aspirated and the cells are resuspended in 0.5 mL of PBS. Mitotic spreads are prepared by pipeting 100 μL of the fixed cells onto a

15 glass microscope slide and flooding the sample with 1 ml of fix solution. Slides then are air dried, stained with Wrights stain (Sigma, St. Louis, MO) for 1 min, followed by one wash in water and one wash in 50% MeOH. The presence of condensed chromosomes and lack of nuclear en-

20 velope identified mitotic cells. Chk1 inhibitors result in an increase in the number of mitotic cells in the presence of irradiation, thereby demonstrating abrogation of the IR-induced G2 arrest. This checkpoint abrogation results in an enhancement in the activity of CyclinB/-

25 cdc2, which is required for progression of cells into mitosis. Cells treated with IR followed by Chk1 inhibitor thus progress into mitosis with damaged DNA. These experiments confirm the hypothesis that Chk1 is involved in the IR-induced G2.

Example 7Chk1 Inhibitor Is Taken Up by Tumor
Cells in the Presence of Chk1 Activator
in a Xenograft Tumor Model

5 In a xenograft tumor model, nude mice are en-
grafted with HT29 colon carcinoma tumors on the flank and
allowed to grow to 200 mm³. Mice then are treated with
either vehicle, 300 mg/kg Chk1 inhibitor, 20 mg/kg
gemcitabine or coadministered with 300 mg/kg Chk1 inhib-
10 itor and 20 mg/kg gemcitabine two times, three days apart
on Days 1 and 4. Treatment of tumor-bearing mice by co-
administration of Chk1 inhibitor and gemcitabine results
in a four-day growth delay in tumors compared to
gemcitabine alone.

15 To assess the diffusion of Chk1 inhibitors
into tumor tissue, plasma and tissue levels of Chk1 in-
hibitor are measured. Using an Alzet pump, 500 mg/kg Chk1
inhibitor is administered to HT29 tumor-bearing mice in a
continuous delivery system over a 24 h period. Plasma
20 samples are taken, then tumors, kidney, liver, spleen,
and lung are harvested. Time points are collected at 1,
2, 4, 8, and 24 h. Tissues are extracted and levels of
Chk1 inhibitor are quantified. This experiment demon-
strates that a Chk1 inhibitor penetrated into normal and
25 tumor tissue, reaches a level of about 15 μ M in tumor
tissue, and peaks in spleen tissue at 8 h at about 20 μ M.
Thus, Chk1 inhibitors were readily taken up by the pro-
liferating cells and are useful, in conjunction with Chk1
activating chemotherapeutic agents, as therapies for the
30 treatment of proliferative diseases.

Example 8**Dose Response of Tumors Treated
with Chk1 Inhibitors and Gemcitabine**

To determine an efficacious dose of Chk1 inhibitor following gemcitabine treatment and whether the dose-dependent checkpoint abrogation correlated with antitumor activity, a dose response experiment is performed.

Nude mice are engrafted with HT29 tumor cells and tumors allowed to develop for 10 days. The tumors at the start were approximately 100 mm³. Animals were treated with gemcitabine at the MTD (160 mg/kg) followed by Chk1 inhibitor at 50 mg/kg, 200 mg/kg, or 400 mg/kg. Gemcitabine pretreatment time is 32 h in this experiment as determined by a cell-based assay that indicated this timepoint as optimal for this type of tumor. Analysis of tumor volume in each treatment regimen indicated that treatment of HT29 tumor bearing mice with the described therapy slows tumor growth greater than gemcitabine alone, with either 200 mg/kg or 400 mg/kg Chk1 inhibitor plus gemcitabine again showing dose-dependent effects of the Chk1 inhibitor.

Example 9**Assay to Determine Whether
an Agent is a Chk1 Activator**

To determine whether an agent is a Chk1 activator, the phosphorylation state of Chk1 can be measured using phospho-specific antibodies to specific phosphorylation sites on Chk1. Serines 317 and 345 have been shown to be phosphorylated after treatment of cells with ionizing radiation, ultraviolet radiation, hydroxyurea, N-

methylnitrosoguanidine (MNNG), temozolamide and gemcitabine. Liu et al., *Genes Dev.* 14:1448-59, 2000; Zhao et al., *Mol. Cell Biol.* 21:4129-39, 2001; Lopez-Girona et al., *Proc. Natl. Acad. Sci. U. S. A.* 98:11289-94, 2001; Guo et al., *Genes Dev.* 14:2745-56, 2000; Gatei et al., *J. Biol. Chem.* 278:14806-11, 2003; Ng et al., *J Biol Chem.* 279(10):8808-19, 2004; Wang et al., *Natl Acad Sci U S A.* 100(26):15387-92, 2003; Stojic et al., *Genes Dev.* 18(11):1331-44, 2004. These serine sites are phosphorylated by upstream checkpoint kinases, Atm and Atr. Liu et al., *Genes Dev.* 14:1448-59, 2000; Zhao et al. *Mol. Cell Biol.*, 21:4129-39, 2001).

The phosphorylation of these sites in response to a candidate Chk1 activator can be monitored by Western blot or immunohistochemistry of tumor cells. For example, the following procedure can be used to demonstrate that gemcitabine results in Chk1 activation at serine 345 and 317. HT29 cells are treated with 20 μ M gemcitabine for two h. The gemcitabine is washed out of the cell growth media and cells are incubated for 22 additional h. Protein lysates are prepared and separated by an SDS-polyacrylamide gel electrophoresis. Proteins are transferred to PVDF membranes and probed with antisera (Cell Signalling) specific for either phosphorylated serine 317 or 345 (Cell Signalling). Western blots show that gemcitabine treatment of HT29 colon carcinoma cells results in the phosphorylation of both serines 317 and 345.

Example 10

Assay to Monitor Chk1 Activity in Response to a Chk1 Inhibitor

It has been found that phosphorylation of Chk1 at serine 296 is stimulated by treatment of tumor cells

with gemcitabine, and that phosphorylation at this site is inhibited by Chk1 inhibitors. Phosphorylation at this site is not inhibited by wortmannin, which inhibits Atm and Atr. Therefore, the phosphorylation of serine 296 is
5 distinct from phosphorylation at serines 317 and 345. In addition, it has been found that this site is phosphorylated in purified Chk1 preparations, suggesting that the purified enzyme is able to phosphorylate itself or other Chk1 molecules at serine 296. Taken together, these data
10 suggest that phosphorylation at serine 296 is performed by Chk1 itself. Therefore, this approach can be used to monitor Chk1 activity in tumors in response to Chk1 activators. Further, this approach can be used to measure inhibition of Chk1 activation by Chk1 inhibitors.

15 Thus, HT 29 cells are treated with 20 μ M gemcitabine for two h. The gemcitabine is washed out of the cell growth media and cells are incubated for 22 additional h. Protein lysates are prepared and separated by an SDS-polyacrylamide gel electrophoresis. Proteins
20 are transferred to polyvinylidene fluoride (PVDF) membranes and probed with antisera (Cell Signalling) specific for phosphorylated serine 296 (Cell Signalling). Western blot shows that gemcitabine treatment of HT29 colon carcinoma cells results in the phosphorylation of
25 serine 296. Further, HT29 cells treated with selective Chk1 inhibitors for 15 min show no serine 296 phosphorylation. These data suggest that serine 296 phosphorylation is performed by the Chk1 kinase.

Example 11

30 Animal Tumor Models

To test the ability of the Chk1 inhibitors of the invention to enhance the killing of tumors by DNA

damaging agents in mice, xenograft tumor models using colon tumor cell lines are established. 5-fluorouracil (5-FU) or gemcitabine can be used as DNA damaging agents. HT29 and Colo205 (human colon carcinoma) and H460 and Calu-6 (nonsmall cell carcinoma) cells can be used to propagate xenograft tumors in 6-8 week old female thymic Balb/c (nu/nu) mice. Mice are maintained in a laminar airflow cabinet under pathogen-free conditions and fed sterile food and water *ad libitum*. Cell lines are grown to subconfluence in RPMI 1640 media supplemented with 10% FBS, 100 U/mL penicillin, 100 μ g/mL streptomycin, and 1.5 mM L-glutamine in a 5% CO₂ humidified environment. Single cell suspensions are prepared in CMF-PBS, and cell concentration adjusted to 1×10^8 cells/mL. Mice are inoculated subcutaneously (s.c.) on the right flank or right leg with a total of 1×10^7 cells (100 μ L).


Mice are randomized (5-15 mice/group) into four treatment groups and used when tumors reach a volume of 75-100 cm³ (usually 7-11 days post-inoculation). Tumors are measured with vernier calipers and tumor volumes are estimated using the empirically derived formula: tumor volume (cm³) = tumor length (cm) x tumor width (cm) x tumor depth (cm)/3.3. Treatment consists of i) 100 μ L intraperitoneal (i.p) injection of gemcitabine at 160 mg/kg. A delay in tumor growth is observed in the mice treated with gemcitabine. Treatment of mice with both 160 mg/kg gemcitabine in combination with oral administration of Chk1 inhibitors is expected to reduce tumor volumes and prolong life. Tumor size is monitored every other day for the duration of the experiment.

Obviously, many modifications and variations of the invention as hereinbefore set forth can be made without departing from the spirit and scope thereof, and,

therefore, only such limitations should be imposed as are indicated by the appended claims.

WHAT IS CLAIMED IS:

1. A compound having a formula



wherein X¹ is null, -O-, -S-, -CH₂-, or -N(R¹)-;

X² is -O-, -S-, or -N(R¹)-;

Y is O or S; or =Y represents two hydrogen atoms attached to a common carbon atom;

W is selected from the group consisting of heteroaryl, aryl, heterocycloalkyl, cycloalkyl, and C₁₋₆alkyl substituted with a heteroaryl or aryl group, wherein said aryl group W is optionally substituted with one to four substituents represented by R², said heteroaryl group W is optionally substituted with one to four substituents represented by R⁵, and said heterocycloalkyl and cycloalkyl groups W are optionally substituted with one or two C₁₋₆alkyl substituents;

R¹ is selected from the group consisting of hydro, C₁₋₆alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, and aryl;


R² is selected from the group consisting of halo, optionally substituted C₁₋₆alkyl, C₂₋₆alkenyl, OCF₃, NO₂, CN, NC, N(R³)₂, OR³, CO₂R³, C(O)N(R³)₂, C(O)R³, N(R¹)COR³, N(R¹)C(O)OR³, N(R¹)C(O)C₁₋₆alkyleneC(O)R³, N(R¹)C(O)C₁₋₆alkyleneC(O)OR³, N(R¹)C(O)C₁₋₆alkyleneOR³, N(R¹)C(O)C₁₋₆alkyleneNHC(O)OR³, N(R¹)C(O)C₁₋₆alkyleneSO₂NR³, C₁₋₆alkyleneOR³, and SR³;

R³ is selected from the group consisting of hydro, halo, C₁₋₆alkyl, C₂₋₆alkenyl, cycloalkyl, aryl,

heteroaryl, SO_2R^4 , C_{1-6} alkyl substituted with one or more of halo, hydroxy, aryl, heteroaryl, heterocycloalkyl, $\text{N}(\text{R}^4)_2$, and SO_2R^4 , C_{1-6} alkylenearyl, C_{1-6} alkyleneheteroaryl, C_{1-6} alkylene C_{3-8} heterocycloalkyl, C_{1-6} alkylene SO_2 aryl, optionally substituted C_{1-6} alkylene $\text{N}(\text{R}^4)_2$, OCF_3 , C_{1-6} alkylene $\text{N}(\text{R}^4)_3^+$, C_{3-8} heterocycloalkyl, and $\text{CH}(\text{C}_{1-6}$ alkylene $\text{N}(\text{R}^4)_2)_2$, or two R^3 groups are taken together to form an optionally substituted 3- to 8-membered aliphatic ring;

R^4 is selected from the group consisting of hydro, C_{1-6} alkyl, cycloalkyl, aryl, heteroaryl, C_{1-6} alkylenearyl, and $\text{SO}_2\text{C}_{1-6}$ alkyl, or two R^4 groups are taken together to form an optionally substituted 3- to 8-membered ring;

R^5 is selected from the group consisting of C_{1-6} alkyl, C_{2-6} alkynyl, aryl, heterocycloalkyl, $\text{N}(\text{R}^3)_2$, OR^3 , halo, N_3 , CN , C_{1-6} alkylenearyl, C_{1-6} alkylene $\text{N}(\text{R}^3)_2$, $\text{C}(\text{O})\text{R}^3$, $\text{C}(\text{O})\text{OR}^3$, $\text{C}(\text{O})\text{N}(\text{R}^3)_2$, $\text{N}(\text{R}^1)\text{C}(\text{O})\text{R}^3$, $\text{N}(\text{R}^1)\text{C}(\text{O})\text{OR}^3$, and



R^6 is $-\text{C}\equiv\text{C}-\text{R}^7$ or heteroaryl;

R^7 is selected from the group consisting of hydro, C_{1-6} alkyl, aryl, C_{1-6} alkylenearyl, heteroaryl, C_{1-6} alkyleneheteroaryl, and alkoxy;

R^8 , R^9 , and R^{10} , independently, are selected from the group consisting of halo, optionally substituted C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, OCF_3 , CF_3 , NO_2 , CN , NC , $\text{N}(\text{R}^3)_2$, OR^3 , CO_2R^3 , $\text{C}(\text{O})\text{N}(\text{R}^3)_2$, $\text{C}(\text{O})\text{R}^3$, $\text{N}(\text{R}^1)\text{COR}^3$, $\text{N}(\text{R}^1)\text{C}(\text{O})-\text{OR}^3$, $\text{N}(\text{R}^8)\text{C}(\text{O})\text{OR}^3$, $\text{N}(\text{R}^1)\text{C}(\text{O})\text{C}_{1-3}$ alkylene $\text{C}(\text{O})\text{R}^3$, $\text{N}(\text{R}^1)\text{C}(\text{O})-\text{C}_{1-3}$ alkylene $\text{C}(\text{O})\text{OR}^3$, $\text{N}(\text{R}^1)\text{C}(\text{O})\text{C}_{1-3}$ alkylene OR^3 , $\text{N}(\text{R}^1)\text{C}(\text{O})-$

C_{1-3} alkyleneNHC(O)OR³, N(R¹)C(O) C_{1-3} alkyleneSO₂NR³, C_{1-3} alkyleneOR³, and SR³;

and a pharmaceutically acceptable salts, or prodrug, or solvate thereof.

2. The compound of claim 1 wherein


X¹ and X² are -N(H)-;

Y is O or S;


W is heteroaryl containing at least two heteroatoms selected from the group consisting of N, O, and S, said ring optionally substituted with one to four substituents selected from the group consisting of C_{1-6} alkyl, aryl, heteroaryl, N(R³)₂, OR³, C(O)N(R³)₂, CO₂R³, CN, and halo.

3. The compound of claim 2 wherein W is selected from the group consisting of pyridazinyl, pyrimidinyl, pyrazinyl, and triazinyl, optionally substituted with one to four substituents selected from the group consisting of optionally substituted C_{1-6} alkyl, aryl, N(R³)₂, OR³, C(O)OR³, C(O)N(R³)₂, and halo.

4. The compound of claim 2 wherein W is selected from the group consisting of



5. The compound of claim 2 wherein W is selected from the group consisting of




6. The compound of claim 2 wherein W is pyrazinyl.

7. The compound of claim 1 wherein R⁷ is heteroaryl.

8. The compound of claim 7 wherein heteroaryl is pyridyl.

9. The compound of claim 1 wherein R⁶ is heteroaryl selected from the group consisting of



10. The compound of claim 9 wherein hetero-aryl is substituted with C_{1-3} alkyleneN(R⁴)₂.

11. A composition comprising a compound of claim 1 and a pharmaceutically acceptable carrier.

12. A compound selected from the group consisting of: 1-(5-methyl-pyrazin-2-yl)-3-(5-methyl-2-pyridin-3-ylethynyl-phenyl)-urea, 1-(5-methyl-pyrazin-2-yl)-3-(5-methyl-2-pyridin-3-yl-phenyl)-urea, 1-(5-methyl-pyrazin-2-yl)-3-(5-methyl-2-pyridin-4-yl-phenyl)-urea, 1-(5-methyl-pyrazine-2-yl)-3-(2-oxazol-5-yl-phenyl)-urea, 1-(5-methyl-pyrazin-2-yl)-3-(5-methyl-2-thiazol-2-yl-phenyl) urea, 1-[2-(4-dimethylaminomethyl-thiazol-2-yl)-5-methyl-phenyl]-3-(5-methyl-pyrazin-2-yl)-urea, and mixtures thereof.

13. A method of inhibiting checkpoint kinase 1 in a cell comprising a step of contacting the cell with an effective amount of a compound of claim 1.

14. A method of sensitizing cells in an individual undergoing a chemotherapeutic or radiotherapeutic treatment for a medical condition, comprising administering to the individual a therapeutically effective amount of a compound of claim 1 in combination with a chemotherapeutic agent, a radiotherapeutic agent, or a mixture thereof.

15. The method of claim 14 further comprising administering one or more cytokine, lymphokine, growth factor, or other hematopoietic factor.

16. The method of claim 14 wherein the chemotherapeutic agent is selected from the group consisting of an alkylating agent, an antimetabolite, a hormone or antagonist thereof, a radioisotope, an antibody, and mixtures thereof.

17. The method of claim 14 wherein the radio-therapeutic agent is selected from the group consisting of gamma-radiation, X-ray radiation, ultraviolet light, visible light, infrared radiation, and microwave radiation.

18. The method of claim 14 wherein the condition is a cancer selected from the group consisting of a colorectal cancer, a head and neck cancer, a pancreatic cancer, a breast cancer, a gastric cancer, a bladder cancer, a vulvar cancer, a leukemia, a lymphoma, a melanoma, a renal cell carcinoma, an ovarian cancer, a brain tumor, an osteosarcoma, and a lung cancer.

19. The method of claim 14 wherein the condition is a cancer selected from the group consisting of myxoid and round cell carcinomas, a locally advanced tumor, metastatic cancer, Ewing's sarcoma, a cancer metastase, a lymphatic metastase, squamous cell carcinoma, esophageal squamous cell carcinoma, oral carcinoma, multiple myeloma, acute lymphocytic leukemia, acute nonlymphocytic leukemia, chronic lymphocytic leukemia, chronic myelocytic leukemia, hairy cell leukemia, effusion lymphomas (body cavity based lymphomas), thymic lymphoma lung cancer, small cell carcinoma, cutaneous T cell lymphoma, Hodgkin's lymphoma, non-Hodgkin's lymphoma, cancer of the adrenal cortex, ACTH-producing tumors, nonsmall cell cancers, breast cancer, small cell carcinoma, ductal carcinoma, stomach cancer, colon cancer, colorectal cancer, polyps associated with colorectal neoplasia, pancreatic cancer, liver cancer, bladder cancer, primary superficial bladder tumors, invasive transitional cell carcinoma of the bladder, muscle-invasive bladder cancer, prostate cancer, ovarian carcinoma, primary peritoneal epithelial neoplasms, cervical carcinoma, uterine endometrial cancers, vaginal cancer, cancer of the vulva, uterine cancer and solid tumors in the ovarian follicle, testicular cancer, penile cancer, renal cell carcinoma, intrinsic brain tumors, neuroblastoma, astrocytic brain tumors, gliomas, metastatic tumor cell invasion in the central nervous system, osteomas and osteosarcomas, malignant melanoma, tumor progression of human skin keratinocytes, squamous cell cancer, thyroid cancer, retinoblastoma, neuroblastoma, peritoneal effusion, malignant pleural effusion, mesothelioma, Wilms's tumors, gall bladder cancer, trophoblastic neoplasms, hemangiopericytoma, and Kaposi's sarcoma.

20. The method of claim 14 wherein the treatment is administered for an inflammatory condition selected from the group consisting of rheumatoid arthritis, psoriasis, vitiligo, Wegener's granulomatosis, and systemic lupus erythematosus.

21. The method of claim 14 wherein the chemotherapeutic agent comprises gemcitabine, pemetrexed, cisplatin, carboplatin, paclitaxel, or mixtures thereof.

22. A method of inhibiting aberrant cell proliferation comprising contacting a cell population comprising aberrantly proliferating cells with a Chk1 activator to substantially synchronize cell cycle arrest among said aberrantly proliferating cells, and subsequently contacting said cell population with a compound of claim 1 to substantially abrogate said cell cycle arrest.

23. The method of claim 22 wherein said Chk1 activator comprises at least one chemotherapeutic agent.

24. The method of claim 22 wherein said Chk1 activator comprises ionizing or ultraviolet radiation.

25. The method of claim 22 wherein said ionizing radiation is administered in conjunction with a radiosensitizer, a photosensitizer, or a mixture thereof.

26. The method of claim 22 wherein said aberrantly proliferating cells are noncancerous.

27. Use of a compound of claim 1 for the manufacture of a medicament for inhibiting checkpoint kinase 1.

28. Use of a compound of claim 1 for the manufacture of a medicament for sensitizing cells in an individual undergoing a chemotherapeutic or radiotherapeutic treatment for a medical indication.

29. Use of a compound of claim 1 for the manufacture of a medicament for inhibiting aberrant cell proliferation in a cell population comprising aberrantly proliferating cells.

30. Use of a compound of claim 1 for the manufacture of a medicament in a cell for the curative or prophylactic treatment of a condition where inhibition of checkpoint kinase 1 is of a therapeutic benefit.

31. An article of manufacture for human pharmaceutical use comprising:

(a) a pharmaceutical composition comprising a compound of claim 1;

(b) a package insert informing that the composition is useful in the treatment of indications involving aberrant cell proliferation; and

(c) an optional container.

32. An article of manufacture for human pharmaceutical use comprising:

(a) a pharmaceutical composition comprising a compound of claim 1;

(b) a package insert informing that the composition is useful as a chemosensitizer or radiosensitizer in a treatment of an indication related to DNA lesions or DNA replication; and

(c) an optional container.

SEQUENCE LISTING

<110> ICOS Corporation

<120> Compounds Useful for Inhibiting CHK1

<130> 27866/40188A

<150> US 60/585,292

<151> 2004-07-02

<160> 1

<170> PatentIn version 3.3

<210> 1

<211> 16

<212> PRT

<213> artificial

<220>

<223> synthetic peptide

<400> 1

Leu	Tyr	Arg	Ser	Pro	Ser	Met	Pro	Glu	Asn	Leu	Asn	Arg	Arg	Arg	Arg
1				5					10					15	

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.